



Linkages of the dynamics of glaciers and lakes with the climate elements over the Tibetan Plateau



Jian Sun^{a,b,c,*}, Tiancai Zhou^{a,g,h,1}, Miao Liu^a, Youchao Chen^d, Hua Shang^c, Liping Zhu^{e,*},
Arshad Ali Shedayi^{a,f}, Huan Yu^h, Genwei Cheng^d, Guohua Liu^b, Ming Xu^c, Wei Deng^d, Jihui Fan^d,
Xuyang Lu^d, Yukun Shaⁱ

^a Synthesis Research Centre of Chinese Ecosystem Research Network, Key Laboratory of Ecosystem Network Observation and Modelling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China

^b State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China

^c Department of Ecology, Evolution, and Natural Resources, School of Environmental and Biological Sciences, Rutgers University, New Brunswick, New Jersey 08901, USA

^d Institute of Mountain Hazard and Environment, Chinese Academy of Sciences, Chengdu, China

^e Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China

^f Department of Biological Sciences, Karakoram International University, Gilgit, Pakistan

^g University of Chinese Academy of Sciences, Beijing 100049, China

^h Chengdu University of Technology, College of Earth Science, Chengdu 610059, China

ⁱ Australian Bureau of Meteorology, Melbourne, GPO Box 1289, VIC 3001, Australia

ARTICLE INFO

Keywords:

Tibetan Plateau
Lakes and glaciers
Regional climate
Water resource

ABSTRACT

Future climate warming is expected to have a significant effect on the operation of Earth and Ecological systems. A key concern in the future is water resource availability. In regions such as the Tibet Plateau (TP) lakes and glaciers appear to be highly sensitive to climate forcing and variations in the size and extent of these system will have profound socio-economic and environmental consequences in South and Central Asia. Although the variety of glaciers and lake son the TP is a heavily researched and discussed topic the interaction between glaciers/lakes and climate change has not be thoroughly investigated. Here we present, through a review of existing studies and original remote sensing analysis, a reconstruction of changes in the spatial coverage of glaciers and lakes on the TP from 1990 to 2015 along with an analysis of climate data for the same period. The results revealed that these systems responded to changes in both temperature and precipitation but the nature of this response, and the controlling factor, was spatially diverse. During this interval the total number of lakes increased from 868 to 1207, thus a large number of new lakes ($n = 339$) formed. The total water surface area of the lakes increase from 38,823.3 km² in 1990 to 48,793.0 km² in 2015, at a rate of 383.5 km² yr⁻¹. Over this period intensive glacial shrinkage occurred, primarily driven by increasing average temperature, except in the Karakoram Mountains where a slight increase in glacier extent was explained by low and stable temperatures along with increasing precipitation. The expansion of lakes in the central and northeastern TP can, therefore, be explained by a trend of increasing precipitation and the accelerated melting of glaciers associated with rising temperatures, both of which contributed to the enhanced total basin runoff. The shrinkage of lake areas along the Himalayan Mountains is accounted for by low precipitation coupled with high evaporation and limited basin space. The lakes within the Qaidam Basin express a complex pattern of response in association with fluctuating precipitation and strong evaporation. The pattern of shrinking glaciers and expanding lakes indicate that water-cycle processes on the TP have been accelerating during the past 25 years. Under current climates, and future climate change, the shrinkage of glaciers and the enlargement of lakes may be expected to continue to accelerate until a “tipping point” is reached when the meltwater of declining glaciers can no longer sustain the enhanced

* Corresponding authors at: Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences (CAS), 11A, Datun Road, Chaoyang District, Beijing 100101, China.

E-mail addresses: sunjian@igsnr.ac.cn (J. Sun), liumiao@igsnr.ac.cn (M. Liu), hua.shang@rutgers.edu (H. Shang), lpzhu@itpcas.ac.cn (L. Zhu), arshadbio@kiu.edu.pk (A.A. Shedayi), gwcheng@imde.ac.cn (G. Cheng), ghliu@rcees.ac.cn (G. Liu), mingxu@igsnr.ac.cn (M. Xu), dengwei@imde.ac.cn (W. Deng), jhfan@imde.ac.cn (J. Fan), xylu@imde.ac.cn (X. Lu), A.Sha@bom.gov.au (Y. Sha).

¹ This author contributed equally to this work.

lake levels. Such a scenario would have severe socio-economic and ecological consequences for this region making the continued study of water-cycle activity across the TP an urgent priority.

1. Introduction

The Tibetan Plateau (TP), the highest and most extensive highland in the world with an extraordinarily sensitive ecosystem, is considered the amplifier of drastic global climate change (Ding et al., 2006; Song et al., 2014d; Sun et al., 2013a; Sun et al., 2013b; Sun et al., 2016; Zhang et al., 2013d). As the critical components in its ecological system, the dynamics of the lakes (Wu and Zhu, 2008; Zhu et al., 2010b) and glaciers (Kaab et al., 2012; Kang et al., 2009b; Molg et al., 2014) are of considerable significance with regard to regional responses to global climate change (Ding et al., 2006; Song et al., 2014b). The variations of hydrological circulation have remarkable impacts on soil conservation (Herman et al., 2015; Kong and Yu, 2013), vegetation productivity (Kramer et al., 2010; Zhang et al., 2013c), biodiversity (Jacobsen et al., 2012), livestock (Shang et al., 2014), land cover types (Molg et al., 2012), tourism (Wang et al., 2010), and socio-economic development (Sheng et al., 2005). Consequently, a number of studies have focused on the dynamics of lakes and glaciers on the TP (Tang et al., 2013).

Lakes play a pivotal role as regulators, sentinels and integrators of climate change (Li et al., 2014b; Yi et al., 2015). Regarding lakes along the Himalayan Mountains, the area of Yamzhog Yumco and Paiku Co sharply decreased in recent decades (Chu et al., 2012; Ye et al., 2007). The lake level of Yamzhog Yumco increased at a rate of 0.28 m yr⁻¹ from 1996 to 2004, but then dropped rapidly at a rate of -0.5 m yr⁻¹ in the period from 2004 to 2012 (Stoffberg et al., 2017; Yang et al., 2008b; Zhang et al., 2015a). The total area of the Mapam Yumco Basin decreased by 34.16 km² from 1974 to 2013 (Ye et al., 2008a). Similarly, the shrinkage of lakes also occurred in the source region of Yellow River (Wan et al., 2014). Lakes were observed to expand, however, in certain regions. In the 2000s, Qinghai Lake began to expand (Liu and Liu, 2008; Shao et al., 2008; Li et al., 2012), with an elevation rate of 0.11 m yr⁻¹ (2003–2009) (Zhang et al., 2011b). The increased rate of the Nam Co Basin was 1.27 km² yr⁻¹ from 1970 to 2000 (Wu et al., 2007b). Between 1976 and 2010, the area of six closed lakes (Siling Co, Nam Co, Bam Co, Pung Co, Darab Co, and Zige Tangco) expanded by 20.2% in total area, accompanied by a 37.7 Gt increase in total storage (Lei et al., 2013). Additionally, lakes in both Nagqu and Kekexili have been continually expanding (Song et al., 2014b). In summary, lakes on the TP grew from 35,638.1 km² to 41,938.7 km² from the 1970s to 2011 (Song et al., 2013). The mass balance of lakes on the TP gained in the 2000s (Lei et al., 2013; Song et al., 2014a; Zhang et al., 2013a), regardless of the area or number of lakes (Wu et al., 2007a). Further, lakes' expansion accelerated from 2003 to 2009 (Zhang et al., 2011b; Zhang et al., 2011c), but the status of lakes on the TP is characterized by large water quantity and spatiotemporal variability.

Glacial melting is expected to impact the hydrologic cycle on regional and global scales (Yang et al., 2015). The glaciers on the TP act as a water storage tower and provide fresh water to almost 1 billion people in South and East Asia (Xu et al., 2009a). A large number of studies analyzed the spatial-temporal dynamics of glaciers on the TP (Aizen et al., 2007; Wang et al., 2011a), and a marked shrinkage of glaciers was revealed in recent decades (Bolch, 2007; Kong and Pang, 2012). Almost 17% of the Great Himalayan is covered by glaciers (Dyurgerov and Meier, 2004), but 50% of the glaciers in the Greater Himalaya of Zaskar (2001–2007), 84% of the glaciers in the western Himalaya (2001–2007), and 42% of the glaciers in the Karakoram (2001–2006) all exhibit a retreating trend (Bolch et al., 2012). In the west of TP, negative mass balance was found in the Dokriani glaciers (1992–2000) (Dobhal et al., 2008), and glaciers in the Mapam Yumco Basin decreased by 7.53 km² from 1974 to 2003 (Ye et al., 2008a). In

central TP, glaciers retreated in the western Nyainqentanglha Range (Bolch et al., 2010b; Pu et al., 2006; Zhang et al., 2004), where 22% of the glacial mass disappeared from 1977 to 2010 (Shahtahmassebi et al., 2012). In the southeast of TP, the area of the Zhadang Glacier and the Gurenhekou Glacier have been continually reduced in the last decades (Yu et al., 2013), and the Halong Glacier experienced a significant shrinkage in length at a rate of -10 m yr⁻¹ from 1981 to 2006 (Yao et al., 2012).

It is worth noting that numerous studies have explored the driving forces for the variations of lakes and glaciers under climate change (Wang et al., 2013b). The expansion of most lakes across the TP was caused by the accelerated melting of glaciers due to the rising temperature, as the number of glacial lakes increased in the Himalaya (Wang et al., 2011b), the central TP (Wang et al., 2013b), and the TP (Zhang et al., 2015b). Meanwhile, Zhu et al. (2010a) suggested that the increased temperature induced the melting of the glaciers, as well as the increased precipitation and the decreased evaporation, which caused a significant increase in the storage of Nam Co. The area of the Nam Co catchment increased by 37.45 km², while the glacial area decreased by 25.74 km² from 1970 to 2000 (Wu and Zhu, 2008). The decrease of lake evaporation contributed to about 4% of the increased water level in Nam Co (1998–2008) (Ma et al., 2016). Generally, many studies have provided abundant evidence for the dynamics of lakes and glaciers on the TP under climate fluctuations. Lei et al. (2014b) hold that the increased regional precipitation contributed to the significant expansion of closed lakes on the TP during 1990 to 2010, while others argue that the expansion of glacier-fed lakes was mainly caused by the increased temperature that accelerated the shrinkage of glaciers (Huang et al., 2011a; Yao and Yu, 2007; Ye et al., 2007). In addition to the melting glaciers, increasing precipitation and declining evaporation were also found to be crucial to the expansion of lakes in central Tibet (Bian et al., 2010; Zhu et al., 2010a).

Many studies have investigated the hydrology of lakes (Wu et al., 2014; Zhang et al., 2011a) but ignored the glaciers (Lei et al., 2013), or only focused on the lakes and glaciers within a small watershed (Gao et al., 2012a). Increasing temperature and precipitation, as well as the shrinkage of glaciers, could accelerate the water cycle (Cheng and Wu, 2007; Gong et al., 2006; Jin et al., 2009; Kang et al., 2010b), especially in recent decades. Because the climate-driven mechanisms of lake and glacier variations still remain unclear in different regions over the TP (Song et al., 2014b), more comprehensive studies are necessary to clarify the relationships among glaciers, lakes and climate. Therefore, this review seeks to answer the following questions: (1) What is the spatial distribution and variations of lakes and glaciers on the TP from 1990 to 2015? (2) What is the relationship between the dynamics of lakes and glaciers under global climate change scenarios?

2. Data sources and analytical methods

2.1. Glacier and lake database

In this study, a time series of satellite images were obtained from the Geospatial Data Cloud (<http://www.giscloud.cn/>) and the United States Geological Survey (www.usgs.gov/). A total of 903 Landsat scenes of Thematic Mapper (TM), Enhanced Thematic Mapper (ETM+), and Operational Land Imager (OLI) were downloaded in the periods of 1990, 1995, 2000, 2005, 2010, 2013 and 2015 and most of the images had a cloud cover of less than 10%. The spatial distribution of images in 1990 and 2015 are exhibited in Appendix Fig. S1 A, B. The images, mainly from June to October in each research period, were selected

because summer precipitation is the main water source for lakes. Detailed image information appears in Appendix Table 1. Landsat images are characterized by the high spatial resolution and their objectivity, and they have been widely used for acquiring the status of glaciers and lakes globally (Bolch et al., 2010a; Wang et al., 2013b; Zhang et al., 2015b). Extracted images were utilized to monitor the status of the lakes in 1990, 1995, 2005, 2010, 2013, and 2015, and the glacial status in 1990 and 2015. Meanwhile, visual interpretation and manual vectorization were conducted by experienced remote sensing experts to re-correct the inaccurate water and glacier surface areas, which were checked using Google Earth images. Finally, the rivers and reservoirs were excluded by consulting lake experts, and a database of lakes and glaciers in the study area was established (Fig. 1).

2.2. Climate data

Regional climate dynamics are important driving factors for glacier and lake variability. The weather station data of temperature, precipitation, and evaporation were collected from the China Meteorological Administration (<http://cdc.cma.gov.cn>) and were processed to obtain the annual mean temperature (AMT), the annual mean precipitation (AMP), and the annual mean evaporation (AME). The spatial interpolation of temperature and precipitation were processed by Anusplin 4.2 (Centre for Resource and Environmental Studies, Australian National University, Canberra), and the Kriging interpolation method was used for the evaporation by ArcGIS 10.2 (ESRI, Inc., Redlands, CA, USA). The spatial changing rates of AMT, AMP, and AME were all calculated by Eq. 1 (Sun et al., 2013a):

$$\beta = \frac{n \sum XY - \sum X \sum Y}{n \sum X^2 - (\sum X)^2} \quad (1)$$

where β is the changing rate of Y , n is the number of years, X is the number of time series (1, 2, 3, ..., n), and Y is the annual average of climate factors in the corresponding year.

2.3. Rule for the partitions on the Tibetan Plateau

The characteristics of precipitation and the dynamics of glaciers on the TP are driven by atmospheric moisture flux: the westerlies (northern TP), Indian monsoon (southern TP), East Asian monsoon (southeastern TP), and the transitions in between (Yao et al., 2013; Yao et al., 2012). The westerlies, Indian monsoon and East Asian monsoon are critical to the climate patterns over the TP (An et al., 2012; Zhisheng et al., 2001). Hence, to systematically and comprehensively reveal the relationship among lakes–glaciers–climate on the TP in different districts, we divided the TP into eight regions (Fig. 8 I A–H) according to the gradient of the large-scale atmospheric circulation that is associated with the westerlies, Indian monsoon and East Asian monsoon.

3. Results and discussion

3.1. Dynamics of lake and glacier

3.1.1. Lakes

The lakes on the TP clearly expanded between 1990 and the 2015 in this study. The lake area shrunk remarkably between 1990 and 1995, then began to continuously expand from 1995 to 2010, and then slowly increased during 2010 to 2015. To better quantify the changes in different-sized lakes, we categorized the lakes into five classes: small (1–2 km²), medium (2–10 km²), big lake (10–50 km²), large lake (50–1,000 km²), and super lake (> 1,000 km²), and detailed information for the number and area of lakes is given in Appendix Table 2. In 1990 and 2015, a total number of 868 and 1,207 lakes with a water surface area larger than 1 km² were detected on the TP, respectively. Small, medium, big, large, and super lakes in 1990 accounted for 26.0%, 36.1%, 21.5%, 15.9%, and 0.5%, respectively, compared to 29.7%, 35.7%, 20.2%, 13.8%, and 0.5% in 2015, respectively (Fig. 2B). The total number of lakes increased markedly by 41.6% from 1990 to 2015, and the biggest change occurred in small lakes with a large number (339) of new lakes formed. The total water surface area of the lakes increased from 38,823.3 km² in 1990 to 48,793 km² in 2015 with an




Fig. 1. The study area.




Fig. 2. Spatial-temporal distribution and variations of lakes on the Tibetan Plateau from 1990 to 2015. Graphs A and B represent the area and number of lakes, respectively; graph C represents the changed portion of lakes from 1990 to 2015; and the 90 represent the year of 1990, 95–1995, 00–2000, 05–2005, 10–2010, 13–2013, 15–2015 in graphs A and B.

increase rate of 20.4% (Fig. 2A). The small, medium, big, large, and super lakes increased by 197.6 km², 614.6 km², 1,102.3 km², 4,533.1 km², and 3,522.2 km², respectively, and the area of large lakes showed the largest increase compared to other sized lakes.

Spatially, Fig. 2C illustrates that the increased lakes are mainly distributed in the Qiangtang Plateau and around the Qinghai Lake. The lake with the greatest expansion was Siling Co, which increased by 647.5 km² (34.8%) from 1990 to 2015 (Appendix Fig. S2). A similar expansion trend of lakes was also found in Tibet (Kleinherenbrink et al., 2015), where a rapid enlargement was exhibited in Siling Co with a growth rate of 26.84 km² yr⁻¹ (2003–2013) (Yi and Zhang, 2015). The water storage of Nam Co also increased by 3.27×10^9 m³, with a total of 2.06 m rising in the lake level (2000–2009) (Zhang et al., 2013c) and a rate of 2.71 km² yr⁻¹ enlargement in the lake area (1976–2009) (Zhang et al., 2011a). Similarly, the area of the lakes in the River Source Region increased by 6,866 ha from the early 1990 to 2004 (Huang et al., 2011b). Qinghai Lake also decreased in lake area between the 1970s and the early 2000s (Feng and Li, 2006; Shao et al., 2008; Shen and Kuang, 2003), while steadily increasing after 2005 (Liu et al., 2013a; Li

et al., 2012). The lakes along the Himalayan Mountains and the Gangdisi Mountains, however, showed a stable unchanging tendency or even a decreasing trend, especially for the area of Yamzhog Yumco, whose lake area decreased by 31.50 km² from 1990 to 2015. Similar results for lakes were also noted in previous studies (Wan et al., 2014; Zhang et al., 2013a). The decreasing rates of 0.40 m yr⁻¹, 0.04 m yr⁻¹, and 0.03 m yr⁻¹ were found in three lakes (Yamzhog Yumco, Peiku Co, Puma Yumco, respectively) in the southwest of the TP (2003–2009) (Zhang et al., 2011d). In addition, the area of lakes in the Hoh Xil region (Fang et al., 2016; Phan et al., 2012) and Qaidam Basin fluctuated in recent years.

The dynamic area data for 78 lakes on the TP was collected from 1990 to 2010 (Fang et al., 2016; Song et al., 2013; Yan and Zheng, 2015) to verify our conclusion. Similar to our results, most of the expanded lakes were distributed in the central TP, and the Har Lake, Qinghai Lake, Eling Lake, and Rige Co in Qinghai province also increased (Fig. 3E–H). The total area of the 53 lakes increased from 20,784.05 km² (1990) to 23,493.37 km² (2010) by 13.04%. The Siling Co expanded the most, with an increasing rate of 29.41 km² yr⁻¹




Fig. 3. Spatial distribution and variations of lakes (km² yr⁻¹) from 1990 to 2010 on the Tibetan Plateau collected from previous studies. Graphs A, B, C and D represent the decreased lakes around the Mapam Yumco, the Yamzhog Yumco, the center and northeastern of the TP, respectively; graphs E–H represent the increased lakes mainly distributed in the center of the TP.

(Fig. 3H). Twenty-one lakes tended to decrease, however, including the lakes in the Himalayan Mountains (Fig. 3A, B, Yamzhog Yumco, Lhanag-tso Lake, and Ngangla Ring Co with decrease rates of $-1.80 \text{ km}^2 \text{ yr}^{-1}$, $-0.54 \text{ km}^2 \text{ yr}^{-1}$, and $-0.39 \text{ km}^2 \text{ yr}^{-1}$, respectively), the left of Bayan Har Mountains (Nianji Co at a rate of $-0.08 \text{ km}^2 \text{ yr}^{-1}$) (Fig. 3D), the Qaidam Basin (Fig. 3D), and a few lakes in the central TP (Sun Lake, Cuona Lake, and Laorite Co with corresponding rates of $-0.03 \text{ km}^2 \text{ yr}^{-1}$, $-0.18 \text{ km}^2 \text{ yr}^{-1}$, and $-0.44 \text{ km}^2 \text{ yr}^{-1}$) (Fig. 3C). In general, the water level of lakes on the TP have risen in recent decades (Liu et al., 2009a; Wang et al., 2013a; Yan and Zheng, 2015), as the area and number of lakes have expanded coherently and significantly (Lei et al., 2014b). The lakes on the TP are characterized by large quantity and spatio-temporal variability.

3.1.2. Glaciers

Glaciers on the TP have been experiencing a dramatic melting and shrinkage in the past 25 years. The status of glaciers is closely related to the real-time regional climate in the highlands; hence, we compared many images to acquire the minimum area of glaciers in each period. Meanwhile, to accurately reflect the dynamics of glaciers on the TP, we not only analyzed the spatial changes in the glacial region in this study but also collected the quantitative dynamics of glaciers from previous studies. Fig. 4 illustrates that glacial retreat was the dominant trend on the TP from 1990 to 2015. A large area and number of glaciers was retreating in the southeast (the Hengduan Mountains), the southwest

(Gangdise Mountains and Himalaya Mountains), the center (the Nyainqentanglha Mountains and Tanggula Mountains), and the north-east (the Qilian Mountains) of the TP. The extremely sensitive glaciers are mainly distributed along the west Gangdise Mountains to the east Himalaya Mountains, and most of the glaciers have been shrinking continuously with a few small mountain glaciers disappearing completely in the Himalaya main ridge (Appendix Fig. S3). The glaciers in the Karakoram Mountains exhibited a slightly positive mass balance, however, and similar results were revealed in previous studies (Bolch and Stoffel, 2012; Kang et al., 2015; Yao et al., 2012). In general, most of the glaciers on the TP showed a negative mass balance (Kang et al., 2009a; Kehrwald et al., 2008; Pu et al., 2008; Ye et al., 2006), and an accelerating decreasing trend was observed in recent decades (Yao et al., 2012).

To better demonstrate our results, the dynamics of glaciers over the TP from previous studies were gathered (Appendix Table 3). The glaciers around the TP exhibited a drastic ablation trend in recent years (Fig. 5) (Li et al., 2008; Zhang et al., 2012c). The extremely sensitive glacial systems were mainly distributed in the edge-region of the TP (changes in area were approximately -10% to -20% , Fig. 5), and the relatively steady glacial systems were primarily found in the west Kunlun Mountains and around the Karakoram Mountains (changes in area were almost -5% , Fig. 5) (Wang et al., 2008b).

For the northwest of the TP, glaciers in the eastern Pamir decreased at an annual rate of $-0.24\% \text{ yr}^{-1}$ (1963–2009) (Zhang et al., 2016). The




Fig. 4. Spatial distribution of the changed glaciers on the Tibetan Plateau between 1990 and 2015.




Fig. 5. Glacial dynamic on the TP (Appendix Table 3). The size of the blue dots denotes the magnitude of glacier area change, and the number with the dot shows the percentage of glacier area change and its time period.

loss ratio of glacial mass in the Muztag Ata and Konggur mountains was -11.2 m yr^{-1} (1990–1999) (Hou et al., 2011; Shangguan et al., 2006) and was $-0.20 \pm 0.01 \text{ m yr}^{-1}$ (1968–1999) in the north bank of the Bangong Co Basin (Wei et al., 2015). The total glacial area decreased by 2.63 km^2 in the Xiongcaigangri region (1968–2013) (Li et al., 2015) and 16 km^2 (1991–2013) in the Depuchangdake region (Li et al., 2016).

In the west of the Himalaya Mountains, the Kangwure Glacier had a significant mass deficit, with a 34.2% glacial area loss since the 1970s (Ma et al., 2010). The changes in average thickness of the Kangwure glacier (2007–2010) and Naimona'nyi glacier (northern branch, 2008–2013) were -2.70 m yr^{-1} and -1.11 m yr^{-1} , respectively (Tian et al., 2014). In the middle of the Himalaya Mountains, the glaciers in the Mt. Qomolangma region decreased by 1.44 km^2 (1992–2000), 1.14 km^2 (2000–2003), and 0.52 km^2 (2003–2008), respectively (Ye et al., 2009a). In the east of the Himalaya Mountains, the total glaciers in the Mapam Basin monotonically decreased, and the ablation rates of glaciers were $-0.20 \text{ km}^2 \text{ yr}^{-1}$ (1973–1990), $-0.32 \text{ km}^2 \text{ yr}^{-1}$ (1990–1999), and $-0.36 \text{ km}^2 \text{ yr}^{-1}$ (1999–2003), with an accelerating ablation trend (Bruehlheide et al., 2011; Ye et al., 2008b).

As for the center of the TP, glacial area in the Geladandong Mountain decreased 128.85 km^2 from 1973 to 2011 (Zhang et al., 2013b), the change in average thickness of Gurenhekou glacier was -3.82 m yr^{-1} between 2007 and 2011 (Tian et al., 2014), and the glacial runoff in the source region of Yangtze River contributed 11.0% of the total river runoff from 1961 to 2000 (Liu et al., 2009b). The glaciers in Nyainqentanglha range (area $> 5.0 \text{ km}^2$) shrank by 22.1% (1999–2013), with area change rates of $-1.32\% \text{ yr}^{-1}$ (1975–1999) and $-1.29\% \text{ yr}^{-1}$ (1999–2013) (Ji et al., 2016). The decreasing area of Zhadang Glacier and Gurenhekou Glacier were 0.64 km^2 and 0.23 km^2 , respectively, from 1970 to 2007 (Yu et al., 2013).

In the southeast of the TP, the area of the Hailuogou Glacier decreased by 0.92 km^2 (1966–2007) (Liu et al., 2010a). The rates of shrinkage in the Hailuogou Glacier tongue and Baishui Glacier No. 1 were -6.16 m yr^{-1} (1990–1997) and -2.70 m yr^{-1} (1997–2004) water equivalent (Zongxing et al., 2010), respectively. The accumulated mass balance in the Hailuogou Glacier was -10.83 m water equivalent

(1959–2003) (Yuanqing et al., 2008), with a retreating rate of $-1.1 \pm 0.4 \text{ m yr}^{-1}$ from 1966 to 2009 (Zhang et al., 2010). Additionally, the 74 glaciers in Gongga Mountains shrank by 29.2 km^2 (1966–2009) (Pan et al., 2012).

The total glacial area in the eastern Qilian Mountains decreased by 1.20 km^2 (1972–2010) (Cao et al., 2014), and the area of the July 1st Glacier decreased by 5% (1956–2002) (Sakai et al., 2005). In the west of Qilian Mountains, the elevation level in Laohugou No. 12 glaciers increased by $18.6 \pm 5.4 \text{ m}$ (1957–2007) (Zhang et al., 2012b), and the area of Qiyi glacier declined from 3.08 km^2 to 0.58 km^2 (1990–2011) (Guo et al., 2015). Meanwhile, the ice volume for the Daxue Range (1957–2010) and the Danghenan Range (1966–2010) declined by 22.40% and 25.70%, respectively (Wang et al., 2016). In the east of the Qilian Mountain, the average glacial thickness in the Ningchan River Glacier No. 3 (1972–2009) and the Shuiguan River No. 4 Glacier (1972–2007) shrank by 9.4 m and $15 \pm 8 \text{ m}$, respectively (Li et al., 2010; Liu et al., 2013b).

3.2. Climate trends

A significant ($P < 0.05$) increase of AMT was observed on the TP from 1990 to 2013, with a fluctuating upward trend from -1.05°C (1990) to 0.84°C (2013), and the average AMT was 0.49°C (Fig. 6A). In the period from 1990 to 2013, the AMT in the northwest and south of TP was relatively higher than in the northeast and the center of TP (Appendix Fig. S4 A). The temperature experienced a more rapid rise between 1990 and 1996 compared to other periods, while it decreased sharply from 1996 to 1997. Specifically, the average AMT was -0.47°C in 1990–1995, 1.06°C in 1996–2000, 0.42°C in 2001–2005, and 0.88°C in 2006–2013. Spatially, the relatively higher AMT appeared first in the southeast and northwest, then in the northeast and northwest, and the lowest value appeared in the middle region. Most areas of the TP experienced an increase in temperature, primarily distributed along the east of Himalaya Mountains and in the center and the east of the TP (Fig. 6D). Simultaneously, the decreasing temperature mainly manifested in the north and southeast of the TP, especially around the




Fig. 6. Spatial patterns and variations of climate factors on the Tibetan Plateau from 1990 to 2013. Graphs A, B and C represent the dynamic of annual mean temperature (AMT), annual mean precipitation (AMP) and annual mean evaporation (AME), respectively; graphs D, E and F represent the spatial change rate of AMT, AMP and AME, respectively, on the TP.

Karakoram Mountains and the northwest of Sichuan Province (Fig. 6D).

During 1990 to 2013, a significant ($P < 0.05$) increase of AMP was observed on the TP, and the average AMP was 348 mm (Fig. 6B). The increasing trend of the AMP fluctuated with an average increase rate of $12.5 \text{ mm/decade}^{-1}$ (1990–2013), and the averages of AMP were 332 mm in 1990–1995, 340 mm in 1996–2000, 352 mm in 2001–2005, and 362 mm in 2006–2013. Spatially, the average AMP presented a descending gradient from the southeast to the northwest (Appendix Fig. S4 B). Moreover, from 1990 to 2013, an increasing trend in precipitation was observed in most parts of the TP, especially in the central TP (Fig. 6E). The AMP remained relatively stable in the Qaidam Basin and declined in the south of TP, however, especially in the Hengduan Mountains and the east of Himalayan Mountains (Fig. 6E).

In general, the AME on the TP showed a fluctuating trend in the period from 1990 to 2013, with a slightly decreasing trend at a rate of 10.5 mm yr^{-1} (1990–2013). A similar phenomenon was found in previous studies, where the AME decreased at a rate of $13.1 \text{ mm decade}^{-1}$ from 1961 to 2000 on the TP (Chen et al., 2006) and 12 mm yr^{-1} in the Namco basin (1998–2008) (Ma et al., 2016). The AME decreased from 1990 to 2003 and then increased from 2004 to 2013. It was observed that the AME from the east to the west of the TP gradually increased, and the west of TP accounted for a larger weight of total evaporation. In detail, higher AME was mainly distributed in the Ngari Prefecture, along the Himalayan Mountains and in the Qaidam Basin (Appendix Fig. S4 C). From 1990 to 2013, a decrease in evaporation was observed in the central TP from north to south, whereas an increasing tendency of AME was observed in the east Pamir Plateau, around the Mapam Yumco, Yamzhog Yumco and Cuorendejia Lake (Fig. 6F).

Overall, in recent decades, climate change on the TP has been characterized by rising temperatures (Wang et al., 2008a) that increased significantly in the eastern and central TP (Du and Ma, 2004; Xu et al., 2008; You et al., 2009). The AMT increased 1.89°C in this study period (1990–2013) on the TP, and the temperature rose 0.3°C per decade in most areas with an altitude above 4,000 m over the past three decades (Xu et al., 2009b). The precipitation was characterized by a rising trend in the central and eastern TP (Rangwala et al., 2009; Xu et al., 2008), but remained relatively stable in the Qaidam Basin and significantly decreased in the south. Meanwhile, no obvious variation was found in the evaporation on the TP during the study period. Slightly decreasing evaporation around the center of TP (Zhang et al., 2009; Zhang et al., 2007) or a relatively stable trend in the southwest (Liao et al., 2013) to the center–north of TP (Huang et al., 2011b) were revealed.

3.3. Dynamics of glaciers respond to the climate change in typical regions

To systematically understand the relationship between glaciers and climate fluctuations on the TP, eight typical regions (Fig. 7 I, A–H) were selected according to the gradient of the large-scale atmospheric circulation as well as their geographic features and dynamic representation of the glaciers. Data on the cumulative mass balance (CMB) of glaciers (Fig. 7 II, A–H) and the AMT (Fig. 7 III, A–H) in each region (Kang et al., 2015; Pan et al., 2012; Wang et al., 2015; Yao et al., 2012) were also collected. Simultaneously, the AMP (Fig. 7 IV A–H) was calculated by the extent of each region A–H in Fig. 7 I.

Significantly increased precipitation ($P = 0.050$, Fig. 7 IV A) and a relatively stable temperature trend (Fig. 7 III A) resulted in a slightly positive mass balance of the Muztag Ata Glacier in the Karakoram Mountains (Fig. 7 I region A), with a 298 mm yr^{-1} increase of CMB between 2006 and 2010 (Fig. 7 II A). More than 50% of the 42 glaciers showed a stable or slightly increasing trend with a mean rate of $8 \pm 12 \text{ m yr}^{-1}$ in the Karakoram ranges (2000–2008) (Scherler et al., 2011), and a total mass gain ($0.37 \pm 0.25 \text{ m yr}^{-1}$) in the glaciers was observed in the north of TP (Neckel et al., 2014). The results showed that the average AMT was -0.59°C from 1990 to 2013 in region A (Fig. 7 I), which was favorable for glaciers to remain stable, especially in the Karakoram Mountains with an obvious decrease of AMT (Fig. 6D). The increased precipitation due to the strengthening of westerlies (Lei et al., 2014a; Li and Wang, 2003; Zhao et al., 2012) presumably keeps the glaciers with a high albedo, which delays the melting of the glaciers (Lei et al., 2014a) in the Karakoram Mountains to a certain extent. More research is needed, however, to reveal why the glaciers in the Pamirs Plateau (decreased) and its surrounding glaciers in the Karakoram Mountains (slightly increased) presented different trends.

The Naimona'nyi Glacier (Fig. 7 I region B) exhibited a negative mass balance as a result of the increasing temperature, with an average CMB of $-2,139 \text{ mm}$ from 2005 to 2010 (Fig. 7 II B). A significant increase in temperature ($P < 0.0005$) and a slight increase in precipitation (Fig. 7 III B, Fig. 7 IV B) were found from 1990 to 2010. The high average AMT (0.92°C) from 2000 to 2008 may account for the rapid retreat (-60 m yr^{-1}) of glaciers in the western Himalayas (Scherler et al., 2011). In addition to the contributions from the increasing temperature, the strong average AME (Fig. S4 C) and low average AMP (134 mm) also had negative impacts on the glacial CMB.

Similar variations in CMB and climate variables were observed in region C (Kangwure Glacier, with an average CMB of $-9,508 \text{ mm}$, 1992–2010, Fig. 7 II C) and region H (Qiyi Glacier, with an average CMB of $-7,939 \text{ mm}$, 1989–2010, Fig. 7 II H) in Fig. 7 I. Both regions




Fig. 7. Dynamics of a typical glacier and its climate characteristics in different regions on the TP. The red triangles in Fig. 7 I A–H represent the distribution of glaciers: A–the Muztag Ata Glacier, B–the Naimona'nyi Glacier, C–the Kangwure Glacier, D–the Zhang glaciers, E–the Xiaodongkemadi Glacier, F–the Parlun NO. 12 glacier, G–74 glacier in the Gongga Mountain and H–the Qiyi Glacier; and Fig. 7 II A–H, Fig. 7 III A–H and Fig. 7 IV A–H represent the cumulative mass balance (CMB), annual mean temperature (AMT) and annual mean precipitation (AMP) of glaciers in these regions, respectively.

showed a significantly increasing trend in temperature ($P < 0.0005$) (Fig. 7 III C, H). Recent studies have illustrated the clearly negative trend in the glacial CMB along the Himalayas (Bolch et al., 2012; Scherler et al., 2011). In detail, the mass balance of the Chhota Shigri Glacier was approximately -1 m in the period from 2003 to 2006 (Wagnon et al., 2007); the four glaciers on the Kangri Karpo Mountains retreated 15–19 m from 2006 to 2007 (Yang et al., 2008a); and in the Mt. Qomolangma Region, the glaciers retreated $0.12 \text{ km}^2 \text{ yr}^{-1}$ between 1992 and 2000, $0.22 \text{ km}^2 \text{ yr}^{-1}$ from 2000 to 2003, and $0.07 \text{ km}^2 \text{ yr}^{-1}$ during 2003 to 2008 (Ye et al., 2009b). The increase in temperature (Yao et al., 2012) and the high average AMT (0.47°C) gave rise to the glacial shrinkage in the central Himalayas. In addition, a slight decreasing trend of precipitation was exhibited in the Himalayas (1979–2010) (Yao et al., 2012) due to the weakening Indian monsoon (Thompson and Mountain, 2006; Yao et al., 2012), which has a negative effect on the glacial CMB. Additionally, the glaciers along the Himalayan Mountains have been melting at an accelerated rate in recent years (Berthier et al., 2007; Kehrwald et al., 2008b; Yao et al., 2012). As for region H, the total glacial area in the Su–lo Mountains decreased by 34.7 km^2 (1966–2000) due to the increasing temperature (Wang et al., 2008c). The negative mass balance of $-0.77 \pm 0.35 \text{ m yr}^{-1}$ from 2003 to 2009 (Neckel et al., 2014) in the Qilian Mountains was dominated by the rapidly increasing temperature (Cao et al., 2014; Wang et al., 2016) and the high average AMT (0.49°C), especially the significantly increased summer mean temperature (Liu et al., 2013b). Thus, the increasing temperature (Fig. 6D) in the northeast reduced the albedo of glaciers, by inducing significant glacial melting (Wu and Zhu, 2008).

The results of this study demonstrated that the increasing temperature accounted for the intensive glacial shrinkage in the Nyainqentanglha Mountains, and the change in CMB of the Zhang glaciers (Fig. 7 I region D, 2005–2012) was $-1,132 \text{ mm yr}^{-1}$ (Fig. 7 II D). The climate in this region was characterized by sharply increasing temperatures after 2004 and fluctuating precipitation from 1990 to 2010 (Fig. 7 III D, IV D). The increasing temperature melted the glaciers on the eastern Nyainqentanglha (Ji et al., 2016), and approximately $13.93 \times 10^6 \text{ m}^3$ water volume was melting from the glaciers in Qugaqie basin (2006–2011), which accounted for 15% of the total basin runoff (Li et al., 2014a). The warming temperature (Fig. 6D) in the center of TP accelerated the retreat of the glaciers.

The change in mass balance for the Xiaodongkemadi Glacier (1989–2011, Fig. 7 II E) and the Da Dongkemadi Glacier (2000–2012) (Wang et al., 2015) was -235 mm yr^{-1} and -242 mm yr^{-1} , respectively (Fig. 7 I region E). The mass balance in Xiaodongkemadi Glacier decreased by an average of 6.15 m (1969–2000) (Xing et al., 2010), with the average mass balance of approximately -136 mm yr^{-1} (1955–2008) (Gao et al., 2012a). Generally, the average level of the five glaciers in the Dongkemadi Glaciers increased from $5,421.6 \text{ m}$ in 1996 to $5,593.4 \text{ m}$ in 2006 (Huang et al., 2013), and thinned 7.34 m from 1955 to 2008 based on the model results (Gao et al., 2012a). A strongly significant ($P < 0.0001$) increase in temperature (Fig. 7 III E) with $0.3^\circ\text{C } 10\text{yr}^{-1}$ in the Dongkemadi River Basin was the main driving force of the glacial ablation, and the glacial meltwater accounted for 66% of the increased runoff (1955–2008) (Gao et al., 2012a).

Many glaciers in region F showed dramatic shrinkage (Fig. 7 I). The average glacial reduction rate exceeded 80 m yr^{-1} in the Ata Glacier from 2005 to 2006 (Yao et al., 2012), and a rapid glacial retreat was found in Parlun No. 94 (-0.9 m yr^{-1} , 2005–2010) (Neckel et al., 2014), Parlun No. 10 glacier (-10.2 m yr^{-1}), and Parlun No. 390 glacier (-5.5 m yr^{-1}) from 2006 to 2008 (Song et al., 2015). The high value of AMT (4.44°C) and significantly ($P < 0.001$) increased temperature ($0.359^\circ\text{C decade}^{-1}$) (Fig. 7 III F) resulted in the most negative mass balance of glaciers in the Parlun No. 12, with a CMB of $-7,163 \text{ mm}$ between 2005 and 2010 (Fig. 7 II F). This was consistent with the study of the Baishui Glacier No. 1 that illustrated the mean temperature of the active-layer increased by 0.24°C in 0.5 – 8.5 m depth of the glaciers during 1982 to 2009 (Wang et al., 2014). Additionally, the glacier area

on the eastern Nyainqentanglha Range decreased by almost 40.53% (1975–2013), which was mainly due to the rising temperature rate of $0.34^\circ\text{C } 10\text{yr}^{-1}$ (Ji et al., 2016). Meanwhile, the distinctly decreased precipitation (Fig. 6E) probably accelerated the glacial shrinkage to a certain extent.

An obvious negative CMB was found in region G (Fig. 7 I). The Hailuoguo Glacier retreated 0.47 m from 1990 to 1997 and 0.42 m from 1952 to 2009 (Li et al., 2013; Xie et al., 2010b; Zhang et al., 2012a). The continuously increasing temperature ($P < 0.01$, Fig. 7 III G) and the fluctuating decreasing precipitation (Fig. 7 IV G) have resulted in continuous shrinkage of the glaciers in the Gongga Mountain region (region G), where the glacial area decreased by 14.4 km^2 from 1989 to 2009 (Fig. 7 II G). The high AMT (4.39°C) and the significantly increasing temperature should be responsible for the shrinkage of the glaciers in Gongga Mountains, with an increasing rate of $0.27^\circ\text{C yr}^{-1}$ from 1990 to 2009 (Zhang et al., 2012a). Similarly, rising temperatures, with a trend of $0.28^\circ\text{C decade}^{-1}$ (1988–2005) (Liu et al., 2010a), have had a significant effect on the increasing glacial retreat in the Hailuoguo Glacier (Duan et al., 2013).

Overall, the increasing temperature was the dominant factor for the shrinkage of the glaciers on the TP (Liu and Chen, 2000; Qin et al., 2009; Yao et al., 2012), and the warming rate ranged from 0.16°C to 0.32°C per decade in winter. Approximately 41.2% of the glaciers were highly vulnerable from 1961 to 2007 due to their high sensitivity to climate change (Yang et al., 2015). The shrinkage of the glacial area in the Ertix River Basin, Junggar Basin, Yarkant River Basin, Dang River Basin, Shule River Basin, and Shiyang River Basin was 8%, 6%, 5%, 3%, 4%, and 20% (Kang et al., 2010a), respectively, in the period from 1960 to 2000. The loss of the glaciers was more serious than expected due to climate warming, especially in the Himalayan Mountains (Ma et al., 2010). The retreating rates of the glaciers were much larger after 1990 than in the 1980s due to the warming climate (Ye et al., 2007). The model results indicated that with temperature increase rates of 0.01, 0.03, and 0.05 K yr^{-1} , the glacial area of China would decrease 14%, 40%, and 60%, respectively, by the end of this century (Xie et al., 2006).

3.4. Hydrobalance mechanisms for typical lakes

First, eight regions of lakes were divided (Fig. 8 I, A–H) to systematically analyze the characteristics of the lakes and climate fluctuations on the TP, according to the gradient of the large-scale atmospheric circulation as well as their geographic features and dynamic characteristics. The dynamics of the lakes' area in the regions of A–H are demonstrated in Fig. 8 II. Accordingly, Fig. 8 III, Fig. 8 IV, and Fig. 8 V represent the variations of climate factors (AMT, AMP and AME) in the eight lake regions from 1990 to 2013, respectively, and the climate factors are calculated by the extent of each region A–H, as shown in Fig. 8 I. Then, to further reveal the relationship between the lakes and climate, we selected eight typical lakes for region A (Aksai Chin Lake), region B (Mapam Yumco), region C (Yamzhog Yumco), region D (Siling Co and Nam Co), region E (Man Co), region F (Cuorendejia Lake), region G (Qinghai Lake), and region H (Dongdabuxun hu) (Fig. 8 I). The basin perimeters and dynamic characteristics of the eight typical lake areas are shown in Appendix Fig. S5 A–I. To summarize, different hydrobalance mechanisms of lakes, which can result in different dynamic patterns, are shown in Fig. 9, and graphs A–H in Fig. 9 show the corresponding mechanisms of the lakes in Appendix Fig. S5 A–H.

The area of the lakes in region A (Fig. 8 I) showed an increasing trend at a rate of $7.8 \text{ km}^2 \text{ yr}^{-1}$ (Fig. 8 II A). The climate in this region was characterized by a slight increase in AMT (Fig. 8 III A), as well as a distinct increase in AMP ($P < 0.05$) (Fig. 8 IV A) and AME ($P = 0.056$) (Fig. 8 V A). From 1990 to 2013, the average of AMT, AMP and AME was -0.52°C , 104 mm and more than $2.15 \times 10^4 \text{ mm}$, respectively. The increasing precipitation was closely related to the strengthening westerlies (Yao et al., 2012c), as the westerlies transported more water




Fig. 8. The dynamic and climatic characteristics of the lakes in different regions on the TP. Fig. 8 I A–H represents the spatial distribution of lake regions that were driven by the atmospheric moisture flux: the westerlies (W), Indian monsoon (IM) and East Asian monsoon (EAM), and transition in between (Transition for W + IM, W + EAM, and IM + EAM); Fig. 8 II A–H represents the dynamic of lake area in regions A–H in Fig. 8 I; Fig. 8 III A–H, Fig. 8 IV A–H, and Fig. 8 V A–H represent the variations of annual mean temperature (AMT), annual mean precipitation (AMP) and annual mean evaporation (AME) in regions A–H in Fig. 8 I, respectively.




Fig. 9. Impact of climate variations in determining dynamic patterns of lakes under the atmospheric moisture flux: the westerlies (W), Indian monsoon (IM) and East Asian monsoon (EAM), and transition in between (Transition for W + IM and IM + EAM). The blue triangles in Fig. 8 I represent the precise location of the typical lakes: A–Aksai Chin Lake, B–Mapam Yumco, C–Yamzhog Yumco, D–Siling Co, E–Man Co, F–Cuorendejia Lake, G–Qinghai Lake and H–Dongdabuxun Hu. Graphs A–H in Fig. 9 represent the hydrobalance mechanism of the typical lakes of Aksai Chin Lake, Mapam Yumco, Yamzhog Yumco, Siling Co and Nam Co, Man Co, Cuorendejia Lake, Qinghai Lake and Dongdabuxun Hu, respectively. The red stars represent significant variation in factors and lake; the "+" and "-" represent the positive and negative variation of factors and lake; the black and red arrows represent the influence factors and the dominant factors for the variations of the lakes.

vapor to the northeast of the TP (Naidu et al., 2009; Yan and Zheng, 2015). In addition, the melting of the glaciers induced by the accelerating increased temperature around the basin (Scherler et al., 2011), together with the increasing precipitation (Fig. 7 IV A), generated a large runoff charging into the lakes. This could be the dominate water source for the expansion of Aksai Chin Lake downstream of the Karakoram Mountains (Fig. 9A).

Differently, a decreasing trend in lake area was found in regions B and C (Fig. 8 I), with decreasing rates of $5.87 \text{ km}^2 \text{ yr}^{-1}$ and $5.42 \text{ km}^2 \text{ yr}^{-1}$, respectively (Fig. 8 II B, C). In both B and C, the climate was characterized by a fluctuating AMT, AMP and AME (Fig. 8 III B, C, Fig. 8 IV B, C, Fig. 8 V B, C), with an average AMT, AMP and AME of 0.60°C and 2.26°C , 183 mm and 433 mm, $2.36 \times 10^4 \text{ mm}$ and $2.14 \times 10^4 \text{ mm}$, respectively. In detail, the Mapam Yumco (in region B, Fig. 8 I) showed an obvious decreasing trend (Song et al., 2014c), with the lake area decreasing at a rate of $1.67 \text{ km}^2 \text{ yr}^{-1}$ (Appendix Fig. S5 I). Additionally, the area of Yamzhog Yumco (in region C, Fig. 8 I) also exhibited a decreasing trend in recent years (Song et al., 2014a; Wang et al., 2013a), with a declining rate of $1.21 \text{ km}^2 \text{ yr}^{-1}$ (Fig. S5 I). Similar shrinkage was investigated in Paiku Co from 1999 to 2010 (Lei et al., 2014b). In fact, a negative mass balance in lakes with areas larger than 10 km^2 (Shen et al., 2014; Song et al., 2013) and an increasing trend of lakes with areas larger than 1 km^2 (Wang et al., 2011b) along the Himalayas were revealed. The high evaporation, low precipitation (Li et al., 2014c), limited space of the basin (Fig. S5 B), and the warm air current of the Indian monsoon (Song et al., 2014c) were the main factors in the shrinkage of Mapam Yumco (Fig. 9B). Previous studies also found that the increasing temperature and decreasing precipitation accounted for the shrinkage of Lhanag-tso, which was next to the Mapam Yumco (Shen et al., 2014). Less precipitation resulted in smaller runoff rates for the Paiku Co and Mapam Yumco basins compared to Nam Co (Biskop et al., 2015). Additionally, low precipitation cannot offset the strong evaporation (Song et al., 2014a; Wang et al., 2013a) and the requirements of the hydroelectric power projects in the Yamzho Yumco (Ye et al., 2007) (Fig. 9C). In particular, the decreasing precipitation and the increasing evaporation led to the sharp reduction of the lakes in Yamzho Yumco after 2004 (Li et al., 2014c). Generally, the AMP of $\sim 200 \text{ mm}$ in lakes along the center of Himalayas to the West Gangdise Mountains made it difficult to offset the strong AME (Song et al., 2014c), which was caused by the warmer Indian monsoon. Previous studies have illustrated that the decreasing precipitation and the particular geographical and geomorphologic environment led to the shrinkage of the lakes in the south of TP (Song et al., 2013; Ye et al., 2007), which was in sharp contrast with the significant expansion of lakes on the central TP (Lei et al., 2013). In fact, the glacial meltwater also played an important role in the water balance (Biskop et al., 2016), and approximately 80% of the increase in runoff was caused by the melting of the snow and ice in the Mt. Qomolangma region. High summer temperatures resulted in the significant increase of the runoff in June (69%), July (35%), and August (14%) in Mt. Qomolangma region (1959–2005) (Liu et al., 2010b).

Generally, the expansion of the lakes in the center of TP was dominated by increasing precipitation, glacial meltwater and decreasing evaporation, which generated the enormous runoff charging into the lakes (Fig. 9D). A dramatic increase in area of the lakes was found in region D (Fig. 8 I), and the area of the lakes increased at a rate of $329.29 \text{ km}^2 \text{ yr}^{-1}$ during 1990 to 2015 (Fig. 8 II D). A significant increase in temperature ($P < 0.01$) (Fig. 8 III D) and precipitation ($P < 0.05$) (Fig. 8 IV D) was observed in region D, with the rates of $0.92^\circ\text{C decade}^{-1}$ and $11.73 \text{ mm decade}^{-1}$ from 1990 to 2013, respectively; and the evaporation exhibited a slightly decreasing trend at a rate of 23.5 mm yr^{-1} (1990–2013) (Fig. 8 V D). Similarly, in the center of TP, an increase in precipitation was observed (Liu et al., 2009a) and modeled (Yang et al., 2011). Coherent growth of the lakes in Siling Co and Nam Co was observed during different periods (Liao et al., 2013), with the increasing rates of $26.64 \text{ km}^2 \text{ yr}^{-1}$ and $2.93 \text{ km}^2 \text{ yr}^{-1}$ from

1990 to 2015, respectively (Appendix Fig. S5 I). The AMT in Nam Co increased at a rate of $0.40^\circ\text{C decade}^{-1}$ in the period from 1976 to 2009 (Zhang et al., 2011a). Additionally, in the Siling Co basin, the precipitation increased at a rate of $17.70 \text{ mm decade}^{-1}$, and the evaporation decreased at a rate of $33.35 \text{ mm decade}^{-1}$ in the hot seasons (1966–2013) (Yi and Zhang, 2015). Quantifying analyses showed that the lake inflows, precipitation, and evaporation contributed 49.5%, 22.1%, and -18.3% , respectively, to the total water storage change from 2003 to 2012 in Siling Co (Zhou et al., 2015). Rainfall–runoff, glacial melting, precipitation, lake percolation, and evaporation accounted for 104.7%, 56.6%, 41.7%, -22.2% , and -80.9% , respectively, of the increased lake level of Nam Co (1980–2010) (Wu et al., 2014). The glacial meltwater that was recharged into the runoff was an important water source (Gao et al., 2012a; Shi et al., 2016; Song et al., 2014d), since the glacial runoff accounted for 60% of the total runoff in the Da Dongkemadi Basin (1955–2008) (Gao et al., 2012a). The ice volume for the Siling Co catchment was approximately 36.37 km^3 (Bian et al., 2010), whereas the glacial area decreased by 34.76 km^2 from 1990 to 2011 (Du et al., 2014). Therefore, the rising temperature ($0.56^\circ\text{C decade}^{-1}$ in Siling Co during summer, 2000–2013) (Yi and Zhang, 2015), as well as the humid climate, resulted in massive glacial melting (Yi et al., 2015) and generated the extensive runoff (Gao et al., 2012b) in spring, autumn, and winter in central Tibet (Liu et al., 2009a). The model results also demonstrate that the glacial mass loss accounted for 11.7% (in Siling Co), 28.7% (in Nam Co), and 11.4% (in Pung Co) of the total rising in lake level between 1999 and 2010 (Lei et al., 2013; Lei et al., 2013).

A steady increase in water area was revealed in region E (Fig. 8 I), with an increase rate of $8.37 \text{ km}^2 \text{ yr}^{-1}$ (Fig. 8 II E). The climate in this region was characterized by increasing temperatures ($P = 0.077$) (Fig. 8 III E) with a rate of $0.56^\circ\text{C decade}^{-1}$, stable precipitation ranging from 596 mm to 816 mm, and slightly enhanced evaporation (Fig. 8 IV E, Fig. 8 V E). Through the stable precipitation and the slightly increased evaporation, the precipitation with an average of $\sim 700 \text{ mm}$ and the glacial melting led to a 2.3 km^2 expansion of Man Co (Fig. 9E). Man Co is located in the southeast of TP with all sides coupled by glacial mountains. Even if the persistent increase in temperature accelerated the glacial melting, however, the geographical and geomorphologic environment (Fig. S5 E) limited the continuous expansion of Man Co. Thus, the area of the lakes was also shaped by their tectonics (Yan and Zheng, 2015).

There was an abrupt increase in the area of the lakes in region F (Fig. 8 I), with a rate of $1.61 \text{ km}^2 \text{ yr}^{-1}$ (Fig. 8 II F). A significant increase in temperature ($P < 0.005$, $0.059^\circ\text{C yr}^{-1}$) and precipitation ($P < 0.001$, 2.77 mm yr^{-1}) (Fig. 8 III F, IV F) and a slight increase in evaporation were observed in this region (Fig. 8 V F). A significant increase in precipitation led to the expansion of most lakes in region F. As a non-glacier-fed lake, the Cuorendejia Lake showed a slight enhancement in evaporation and an increase in temperature, which would be the main reasons for the 20.4 km^2 decrease of this lake from 1990 to 2015 (Fig. 9F). Meanwhile, the lack of glacial meltwater supply could be an important reason for the shrinkage of Cuorendejia Lake, since the glacial meltwater acts as a major water source for the lakes on the TP (Gao et al., 2012a; Song et al., 2014d). For example, Siling Co expanded dramatically, while Lake Co (next to Siling Co) remained stable. As a result, the lack of glacial meltwater in Lake Co rendered the two lakes very different, in spite of their similar climate features (Song et al., 2013). The marked variations between Cuorendejia Lake (decreased) and its surrounding lakes (increased), however, suggested that additional mechanisms may be involved in region F (Fig. 9).

The area of lakes in region G (Fig. 8 I) clearly increased at a rate of $37.09 \text{ km}^2 \text{ yr}^{-1}$ (Fig. 8 II G). A significant increase in temperature ($P < 0.05$) (Fig. 8 III G), precipitation ($P < 0.05$) (Fig. 8 IV G), and evaporation ($P = 0.059$) (Fig. 8 V G) was found in region G, with annual average values of -0.11°C , 432 mm, and $1.48 \times 10^4 \text{ mm}$ from 1990 to 2013, respectively. Qinghai Lake is located in the temperate semi-arid

climate zone, which is mainly supplied by precipitation and runoff. The area of Qinghai Lake increased at a rate of $24.32 \text{ km}^2 \text{ yr}^{-1}$ from 1990 to 2015. The significant increasing precipitation (Zhang et al., 2011c) and rapidly melting glaciers (Cao et al., 2014; Wang et al., 2016) gave rise to the expansion of Qinghai Lake. More than 70 of the inflowing rivers that were distributed around Qinghai Lake, together with the enhanced runoff caused by the increasing precipitation, were the primary supply sources for Qinghai Lake (Yan and Zheng, 2015). Meanwhile, the increasing temperature and precipitation made the climate in Qinghai Lake become much warmer and wetter (Liu et al., 2014), which accelerated the melting of the glaciers (with a melting rate of 300 mm per year in Qiye) around the lake, and the glacial water became an important source for the expansion of Qinghai Lake (Song et al., 2013; Zhang et al., 2011c). The evaporation of the lake was much larger than the precipitation (Cui and Li, 2015; Haginoya et al., 2009; Song et al., 2014d), however, where the annual evaporation of the lake offset 23.42% of the precipitation in the basin (Cui and Li, 2015) and weakened the expansion of Qinghai Lake. Additionally, human activities have also had negative impact on the expansion of Qinghai Lake (Yan and Zheng, 2015).

The area of the lakes in region H (Fig. 8 I) increased slightly at a rate of $3.08 \text{ km}^2 \text{ yr}^{-1}$ (Fig. 8 II H). The AMT and AMP both significantly increased (Fig. 8 III H, IV H), with average increase rates of $1.60^\circ\text{C yr}^{-1}$ and 94 mm yr^{-1} , respectively. The evaporation in this region exhibited a stable trend (Fig. 8 V H), with a strong average evaporation exceeding $1.79 \times 10^4 \text{ mm yr}^{-1}$ that was much higher than the precipitation (Xie et al., 2010a; Li et al., 2007). Although a significant increasing trend was observed in precipitation, the climate of this temperate arid climate zone was primarily controlled by evaporation and the accelerating increased temperature (Xie et al., 2010a; Li et al., 2007; Xu et al., 2006). Relatively low precipitation and high evaporation resulted in the uncertain variations of Dongdabuxun Lake (Fig. 9H). Lakes in the Qaidam Basin experienced intense fluctuations from 1990 to 2015, with many lakes newly forming or disappearing throughout the different periods. The obvious expansion of the lakes occurred under relatively abundant precipitation, while dramatic shrinkage appeared in dry seasons. Moreover, the shrinkage of the salt lakes in the Qaidam basin was probably caused by human exploitation (Wan et al., 2014) to acquire the resources of gypsum, mirabilite, and trona (Yan and Zheng, 2015).

In summary, both the increasing precipitation and accelerated melting of the glaciers enhanced the total basin runoff, which were the dominant drivers of the dramatic expansion of lakes in the center and northeast of the TP. The increasing precipitation directly expanded the lakes in the center and northeast of the TP. In addition, the increasing temperature accelerated glacial melting, and the glacier-fed lakes exhibited a much faster increasing trend than non-glacier-fed lakes (Huang et al., 2016). In this study, 133 new lakes ($1 \text{ km}^2 < \text{area} < 2 \text{ km}^2$) were formed from 1990 to 2015. Similarly, a large number (1099) of new lakes with areas larger than 0.003 km^2 were formed on the TP from 1990 to 2010 (Zhang et al., 2015b). The glacial meltwater charging into the lakes in the form of runoff was an important factor in the expansion of lakes, especially at $\geq 33^\circ\text{N}$ north of the TP (Song et al., 2014d).

4. Limitations of current study

Generally, the majority of the studies have simulated the hydrology of lakes (Wu et al., 2014; Zhang et al., 2011a) and either ignored the glaciers and permafrost (Lei et al., 2013) or overlooked the groundwater flowing into lakes (Zhou et al., 2013). For instance, some research has only modeled the mass balance of runoff and glaciers within a small watershed (Gao et al., 2012a), or just predicted single glacier variations (Shi et al., 2016). To accurately estimate the contributions of different factors on lakes, many parameters of water balance are required (Zhu et al., 2010b), including precipitation, evaporation, runoff, and other sub basins, ground water, exact ice mass loss of glaciers.

Because the in situ meteorological observations for every lake and glacier were insufficient, however, it is difficult to estimate climate change. Thus, diverse national and international cooperation is required for developing “lake–atmosphere–glacier” interaction research on the TP.

Moreover, extensive alpine permafrost underlies the TP (Cheng and Liu, 2008). The in situ monitoring demonstrated that the permafrost is becoming thinner and less contiguous due to global warming, and the groundwater from the shrinkage of permafrost could facilitate the expansion of lakes (Liao et al., 2013). This requires further research. Additionally, extracting the surface area of lakes might inevitably induce more or less misestimating. Errors might also occur when simplifying statistical geometry of the lake pixels to obtain the area of lakes as different sensors possess different image resolutions, such as $\sim 1 \text{ km}$ or $\sim 30 \text{ m}$ pixel sizes for MODIS and LANDSAT images, respectively. In addition, the periods of images should be chosen within a constant season window for the area of lakes in order to get the largest extent or maintain relatively stable data.

5. Conclusions and implications

We concluded that the mass balance of glaciers on the TP showed that glaciers are experiencing a rapid retreat process, and the dominant reason for the shrinkage of the glaciers was the increasing temperature. In contrast, a positive mass balance of glaciers around the Karakoram Mountains was observed, where the glaciers exhibited a stable or slowly increasing trend due to the relatively stable temperatures and significantly increased precipitation induced by the strengthening of the westerlies.

Both the area and number of lakes have expanded coherently and significantly on the TP from 1990 to 2015. Spatially, the increasing precipitation and accelerating glacier melting enhanced the total basin runoff, which together formed the dominant drivers for the dramatic expansion of lakes in the center and northeast of the TP. Relatively low precipitation, increased temperature, and geographical and geomorphologic limits (lack of inflowing rivers within basin) led to the shrunken or relatively unchanged lakes along the Himalayan Mountains. Uncertain variations in precipitation, strong evaporation and human activities accounted for the fluctuating changes in lakes in the Qaidam Basin. Therefore, different climate factors varied in hydrobalance mechanisms and generated dynamic patterns of lakes on the TP.

This review enhances our scientific understanding of the responses of water resources and the aquatic ecosystems to climate change. The mountain ranges on the TP function as water towers for the Pacific and Indian Ocean water systems, which have profound impacts on water and ecological security. Under the present climate conditions, it is expected that the lakes on the TP will continually expand for a certain period until they reach a “tipping point,” when the meltwater of declining glaciers cannot offset the demand for water resources. In this context, severe consequences of limited water resources and many unexpected climate effects, which would seriously threaten the human livelihoods and wellbeing in the South and East Asia, might occur. Therefore, future research should devote more attention to the linkages of lakes and glaciers on the TP, which is critical for developing scientific strategies to control or reduce the potential threats.

Acknowledgments

This research was funded by the National Key Research and Development Program of China (Grant No. 2016YFC0502102, 2016YFC0502002, 2016YFC0501803 and 2016YFC0501802), and the National Science Foundation of China (Grant No. 41671016 and 41501057), West Light Foundation of the Chinese Academy of Sciences, and Youth Innovation Research Team Project (LENOM2016Q0003), the National Natural Science Foundation of China (No. 91647110).

Appendix A. Supplementary data

Supplementary data for this article can be found online at <https://doi.org/10.1016/j.earscirev.2018.06.012>.

References

- An, Z., Colman, S.M., Zhou, W., Li, X., E.T., Brown, Jull, A.J.T., Cai, Y., Huang, Y., Lu, X., Chang, H., 2012. Interplay between the Westerlies and Asian monsoon recorded in Lake Qinghai sediments since 32 ka. *Scientific Reports* 2 (8), 619.
- Aizen, V.B., Kuzmichenok, V.A., Surazakov, A.B., Aizen, E.M., 2007. Glacier changes in the Tien Shan as determined from topographic and remotely sensed data. *Global and Planetary Change* 56 (3–4), 328–340.
- Berthier, E., Arnaud, Y., Kumar, R., Ahmad, S., Wagnon, P., Chevallier, P., 2007. Remote sensing estimates of glacier mass balances in the Himachal Pradesh (Western Himalaya, India). *Remote Sensing of Environment* 108 (3), 327–338.
- Bian, D., Bian, B., Ba, L.A., Wang, C., Chen, T., 2010. The Response of Water Level of Selin Co to Climate Change during 1975–2008. *Acta Geographica Sinica* 65 (3), 313–319.
- Bishop, S., Maussion, F., Krause, P., Fink, M., 2015. What are the key drivers of regional differences in the water balance on the Tibetan Plateau? *Hydrology and Earth System Sciences Discussions* 12 (4), 4271–4314.
- Bishop, S., Maussion, F., Krause, P., Fink, M., 2016. Differences in the water–balance components of four lakes in the southern–central Tibetan Plateau. *Hydrology and Earth System Sciences* 20 (1), 209–225.
- Bolch, T., 2007. Climate change and glacier retreat in northern Tien Shan (Kazakhstan/Kyrgyzstan) using remote sensing data. *Global and Planetary Change* 56 (1–2), 1–12.
- Bolch, T., Stoffel, M., 2012. The State and Fate of Himalayan Glaciers. *Science* 336 (6079), 310.
- Bolch, T., Menounos, B., Wheate, R., 2010a. Landsat-based inventory of glaciers in western Canada, 1985–2005. *Remote Sensing of Environment* 114 (1), 127–137.
- Bolch, T., Kang, S.C., Yao, T.D., Buchroithner, M.F., Maussion, F., Scherer, D., Huinjes, E., Schneider, C., 2010b. A glacier inventory for the western Nyainqentanglha Range and the Nam Co Basin, Tibet, and glacier changes 1976–2009. *Cryosphere Discussions* 4 (3), 419–433.
- Bolch, T., Kulkarni, A., Kaab, A., Huggel, C., Paul, F., Cogley, J.G., Frey, H., Kargel, J.S., Fujita, K., Scheel, M., Bajracharya, S., Stoffel, M., 2012. The state and fate of Himalayan glaciers. *Science* 336 (6079), 310–314.
- Bruehlheide, H., Bohnke, M., Both, S., Fang, T., Assmann, T., Baruffol, M., Bauhus, J., Buscot, F., Chen, X.Y., Ding, B.Y., Durka, W., Erfmeier, A., Fischer, M., Geissler, C., Guo, D.L., Guo, L.D., Hardtle, W., He, J.S., Hector, A., Krober, W., Kuhn, P., Lang, A.C., Nadrowski, K., Pei, K.Q., Scherer-Lorenzen, M., Shi, X.Z., Scholten, T., Schuldt, A., Trogisch, S., von Oheimb, G., Welk, E., Wirth, C., Wu, Y.T., Yang, X.F., Zeng, X.Q., Zhang, S.R., Zhou, H.Z., Ma, K.P., Schmid, B., 2011. Community assembly during secondary forest succession in a Chinese subtropical forest. *Ecological Monographs* 81 (1), 25–41.
- Cao, B., Pan, B.T., Wang, J., Shangguan, D.H., Wen, Z.L., Qi, W.T., Cui, H., Lu, Y.Y., 2014. Changes in the glacier extent and surface elevation along the Ningchan and Shuiguan river source, eastern Qilian Mountains, China. *Quaternary Research* 81 (3), 531–537.
- Chen, S.B., Liu, Y.F., Thomas, A., 2006. Climatic change on the Tibetan Plateau: Potential Evapotranspiration Trends from 1961–2000. *Climatic Change* 76 (3), 291–319.
- Cheng, G.D., Wu, T.H., 2007. Responses of permafrost to climate change and their environmental significance, Qinghai-Tibet Plateau. *Journal of Geophysical Research Atmospheres* 112 (F2), 93–104.
- Cheng, Z.G., Liu, X.D., 2008. A preliminary study of future permafrost distribution on the Qinghai-Tibet plateau under global warming conditions. *Areal Research & Development* 27 (06), 80–85.
- Chu, D., Qiong, P.U., Zhuoma, L., Zhu, L., Zhang, X., Ciren, P., Yangzongamp, D., Rui, S., 2012. Remote sensing analysis on lake area variations of Yamzho Yumco in Tibetan Plateau over the past 40 a. *Journal of Lake Sciences* 24 (3), 494–502.
- Cui, B.L., Li, X.Y., 2015. Stable isotopes reveal sources of precipitation in the Qinghai Lake Basin of the northeastern Tibetan Plateau. *Science of the Total Environment* 527, 26–37.
- Ding, Y.J., Liu, S.Y., Bai-Sheng, Y.E., Zhao, L., 2006. Climatic Implications on Variations of Lakes in the Cold and Arid Regions of China during the Recent 50 Years. *Journal of Glaciology & Geocryology* 28 (5), 623–632.
- Dobhal, D.P., Gergan, J.T., Thayyen, R.J., 2008. Mass balance studies of the Dokriani Glacier from 1992 to 2000, Garhwal Himalaya, India, Garhwal Himalaya. *India. Bulletin of Glaciological Research* 25, 9–17.
- Du, J., Yang, T.B., He, Y., 2014. Glaciers and lakes changes and climate response in the Selin Co Basin from 1990 to 2011. *Journal of Arid Land Resources & Environment* 28 (12), 88–93.
- Du, J., Ma, Y.C., 2004. Climatic trend of rainfall over Tibetan Plateau from 1971 to 2000. *Acta Geographica Sinica* 59 (3), 375–382.
- Duan, J., Wang, L., Li, L., Sun, Y., 2013. Tree-ring-inferred glacier mass balance variation in southeastern Tibetan Plateau and its linkage with climate variability. *Climate of the Past* 9 (6), 2451–2458.
- Dyrgerov, M.B., Meier, M.F., 2004. *Glaciers and the Changing Earth System: A 2004 Snapshot*, pp. 58.
- Fang, Y., Cheng, W.M., Zhang, Y.C., Wang, N., Zhao, S.M., Zhou, C.H., Chen, X., Bao, A.M., 2016. Changes in inland lakes on the Tibetan Plateau over the past 40 years. *Journal of Geographical Sciences* 26 (4), 415–438.
- Feng, Z.K., Li, X.H., 2006. Remote sensing monitoring study for water area change and lakeshore evolution of Qinghai Lake in last 20 years. *Journal of Palaeogeography* 8 (1), 131–141.
- Gao, H.K., He, X.B., Ye, B.S., Pu, J.C., 2012a. Modeling the runoff and glacier mass balance in a small watershed on the Central Tibetan Plateau, China, from 1955 to 2008. *Hydrological Processes* 26 (11), 1593–1603.
- Gao, T.G., Kang, S.C., Krause, P., Cuo, L., Nepal, S., 2012b. A test of J2000 model in a glacierized catchment in the central Tibetan Plateau. *Environmental Earth Sciences* 65 (6), 1651–1659.
- Gong, T.L., Liu, C.M., Liu, J.S., 2006. Hydrological response of Lhasa River to climate change and permafrost degradation in Xizang. *Acta Geographica Sinica* 61 (5), 519–526.
- Guo, Z.M., Wang, N.L., Wu, H.B., Wu, Y.W., Wu, X.J., Li, Q.L., 2015. Variations in Firn Line Altitude and Firn Zone Area on Qiyi Glacier, Qilian Mountains, Over the Period of 1990 to 2011. *Arctic, Antarctic, and Alpine Research* 47 (2), 293–300.
- Haginoya, S., Fujii, H., Kuwagata, T., Xu, J.Q., Ishigooka, Y., Kang, S.C., Zhang, Y.J., 2009. Air-lake interaction features found in heat and water exchanges over nam co on the Tibetan Plateau. *Sola* 5, 172–175.
- Herman, F., Beyssac, O., Brughelli, M., Lane, S.N., Leprince, S., Adatte, T., Lin, J.Y.Y., Avouac, J.P., Cox, S.C., 2015. Erosion by an Alpine glacier. *Science* 350 (6257), 193–195.
- Hou, L.L., Liu, Y.F., Wang, X.Y., Ma, H.B., He, J.S., Zhang, Y., Yu, C.H., Guan, W.J., Ma, Y.H., 2011. The effects of amyloid-beta(42) oligomer on the proliferation and activation of astrocytes in vitro. *Vitro Cellular & Developmental Biology-Animal* 47 (8), 573–580.
- Huang, L., Liu, J.Y., Shao, Q.Q., Liu, R.G., 2011. Changing inland lakes responding to climate warming in Northeastern Tibetan Plateau. *Climatic Change* 109 (3), 479–502.
- Huang, L., Li, Z., Tian, B.S., Chen, Q., Zhou, J.M., 2013. Monitoring glacier zones and snow/firn line changes in the Qinghai-Tibetan Plateau using C-band SAR imagery. *Remote Sensing of Environment* 137, 17–30.
- Huang, L., Xu, X.L., Zhai, J., Sun, C.Y., 2016. Local background climate determining the dynamics of plateau lakes in China. *Regional Environmental Change* 16 (8), 2457–2470.
- Jacobsen, D., Milner, A.M., Brown, L.E., Dangles, O., 2012. Biodiversity under threat in glacier-fed river systems. *Nature Climate Change* 2 (5), 361–364.
- Ji, Q., Yang, T.B., He, Y., Chen, J., Wang, K., 2016. Glacier changes in the eastern Nyainqentanglha Range of Tibetan Plateau from 1975 to 2013. *Journal of Mountain Science* 13 (4), 682–692.
- Jin, H.J., He, R.X., Cheng, G.D., Wu, Q.B., Wang, S.L., Chang, X.L., 2009. Changes in frozen ground in the Source Area of the Yellow River on the Qinghai-Tibet Plateau, China, and their eco-environmental impacts. *Environmental Research Letters* 4 (4), 549–567.
- Kaab, A., Berthier, E., Nuth, C., Gardelle, J., Arnaud, Y., 2012. Contrasting patterns of early twenty-first-century glacier mass change in the Himalayas. *Nature* 488 (7412), 495–498.
- Kang, S.C., Chen, F., Gao, T.G., Zhang, Y.J., Yang, W., Yu, W.S., Yao, T.D., 2009. Early onset of rainy season suppresses glacier melt: a case study on Zhadang glacier, Tibetan Plateau. *Journal of Glaciology* 55 (192), 755–758.
- Kang, E., Liu, C.H., Xie, Z.C., Li, X., Shen, Y.P., 2010a. Assessment of glacier water resources based on the Glacier Inventory of China. *Annals of Glaciology* 50 (53), 104–110.
- Kang, S.C., Xu, Y.W., You, Q.L., Flügel, W.A., Pepin, N., Yao, T.D., 2010b. Review of climate and cryospheric change in the Tibetan Plateau. *Environmental Research Letters* 5 (1), 75–82.
- Kang, S., Wang, F., Morgenstern, U., Zhang, Y., Grigholm, B., Kaspari, S., Schwikowski, M., Ren, J., Yao, T., Qin, D., Mayewski, P.A., 2015. Dramatic loss of glacier accumulation area on the Tibetan Plateau revealed by ice core tritium and mercury records. *Cryosphere* 9 (3), 1213–1222.
- Kehrwald, N.M., Thompson, L.G., Yao, T., Mosley-Thompson, E., Schotterer, U., Altimov, V., Beer, J., Eikenberg, J., Davis, M.E., 2008. Mass loss on Himalayan glacier endangers water resources. *Geophysical Research Letters* 35 (22), 113–130.
- Kleinherenbrink, M., Lindenberg, R.C., Dittmar, P.G., 2015. Monitoring of lake level changes on the Tibetan Plateau and Tian Shan by retracking Cryosat SARIn waveforms. *Journal of Hydrology* 521, 119–131.
- Kong, Y.L., Pang, Z.H., 2012. Evaluating the sensitivity of glacier rivers to climate change based on hydrograph separation of discharge. *Journal of Hydrology* 434–435, 121–129.
- Kong, B., Yu, H., 2013. Estimation model of soil freeze-thaw erosion in silingco watershed wetland of northern Tibet. *Scientific World Journal* 3, 636521.
- Kramer, A., Herzschrub, U., Mischke, S., Zhang, C.J., 2010. Late glacial vegetation and climate oscillations on the southeastern Tibetan Plateau inferred from the Lake Naleng pollen profile. *Quaternary Research* 73 (2), 324–335.
- Lei, Y.B., Yao, T.D., Bird, B.W., Yang, K., Zhai, J.Q., Sheng, Y.W., 2013. Coherent lake growth on the central Tibetan Plateau since the 1970s: Characterization and attribution. *Journal of Hydrology* 483 (3), 61–67.
- Lei, Y.B., Yang, K., Wang, B., Sheng, Y.W., Bird, B.W., Zhang, G.Q., Tian, L.D., 2014. Response of inland lake dynamics over the Tibetan Plateau to climate change. *Climatic Change* 125 (2), 281–290.
- Li, X.Y., Xu, H.Y., Sun, Y.L., Zhang, D.S., Yang, Z.P., 2007. Lake-level change and water balance analysis at Lake Qinghai, West China during recent decades. *Water Resources Management* 21 (9), 1505–1516.
- Li, X., Cheng, G.D., Jin, H.J., Kang, E., Che, T., Jin, R., Wu, L.Z., Nan, Z.T., Wang, J., Shen, Y.P., 2008. Cryospheric change in China. *Global and Planetary Change* 62 (3–4), 210–218.
- Li, J., Liu, S.Y., Shangguan, D., Zhang, Y., 2010. Identification of ice elevation change of the Shuiguan River No. 4 glacier in the Qilian Mountains China. *Journal of Mountain Science* 7 (4), 375–379.
- Li, X.D., Xiao, J.S., Li, F.X., Xiao, R.X., Xu, W.X., Wang, L., 2012. Remote Sensing

- Monitoring of the Qinghai Lake Based on EOS/MODIS Data in Recent 10 Years. *Journal of Natural Resources* 27 (11), 1962–1970.
- Li, P., Yang, Y.H., Fang, J.Y., 2013. Variations of root and heterotrophic respiration along environmental gradients in Chinese forests. *Journal of Plant Ecology* 6 (5), 358–367.
- Li, B.Q., Yu, Z.B., Liang, Z.M., Acharya, K., 2014a. Hydrologic response of a high altitude glacierized basin in the central Tibetan Plateau. *Global and Planetary Change* 118, 69–84.
- Li, J., Li, Z.W., Ding, X.L., Wang, Q.J., Zhu, J.J., Wang, C.C., 2014b. Investigating mountain glacier motion with the method of SAR intensity-tracking: Removal of topographic effects and analysis of the dynamic patterns. *Earth-Science Reviews* 138, 179–195.
- Li, L., Li, J., Yao, X.J., Luo, J., Huang, Y.S., Feng, Y.Y., 2014c. Changes of the three holy lakes in recent years and quantitative analysis of the influencing factors. *Quaternary International* 349, 339–345.
- Li, Z.G., Fang, H.Y., Tian, L.D., Dai, Y.F., Zong, J.B., 2015. Changes in the glacier extent and surface elevation in Xiongcaigangri region, Southern Karakoram Mountains China. *Quaternary International* 371, 67–75.
- Li, Z.G., Tian, L.D., Wu, H.B., Wang, W.C., Zhang, S.H., Zhang, J.J., Li, X.X., 2016. Changes in glacier extent and surface elevations in the Depuchangdake region of northwestern Tibet, China. *Quaternary Research* 85 (1), 25–33.
- Li, J., Wang, J.X.L., 2003. A modified zonal index and its physical sense. *Geophysical Research Letters* 30 (12).
- Liao, J.J., Shen, G.Z., Li, Y.K., 2013. Lake variations in response to climate change in the Tibetan Plateau in the past 40 years. *International Journal of Digital Earth* 6 (6), 534–549.
- Liu, X.D., Chen, B.D., 2000. Climatic warming in the Tibetan plateau during recent decades. *International Journal of Climatology* 20 (14), 1729–1742.
- Liu, R.A., Liu, Y., 2008. Area changes of Lake Qinghai in the latest 20 years based on remote sensing study. *Journal of Lake Sciences* 20 (1).
- Liu, J.S., Wang, S.Y., Yu, S.M., Yang, D.Q., Zhang, L., 2009a. Climate warming and growth of high-elevation inland lakes on the Tibetan Plateau. *Global and Planetary Change* 67 (3–4), 209–217.
- Liu, S.Y., Zhang, Y., Zhang, Y.S., Ding, Y.J., 2009b. Estimation of glacier runoff and future trends in the Yangtze River source region, China. *Journal of Glaciology* 55 (190), 353–362.
- Liu, Q., Liu, S.Y., Zhang, Y., Wang, X., Zhang, Y.S., Guo, W.Q., Xu, J.L., 2010a. Recent shrinkage and hydrological response of Hailuoguo glacier, a monsoon temperate glacier on the east slope of Mount Gongga, China. *Journal of Glaciology* 56 (196), 215–224.
- Liu, W.G., Ren, J.W., Qin, X., Liu, J.S., Liu, Q., Cui, X.Q., Wang, Y.T., 2010b. Hydrological characteristics of the Rongbuk Glacier catchment in Mt. Qomolangma region in the central Himalayas China. *Journal of Mountain Science* 7 (2), 146–156.
- Liu, B.K., Wei, X.L., Du, Y.E., Feng, S.Q., Mei, D., Li, C.Y., Liu, Y.Z., Zhang, J.K., 2013a. Dynamics of Qinghai Lake area based on environmental mitigation satellite data. *Pratacultural Science* 30 (02), 178–184.
- Liu, Y.S., Qin, X., Zhang, T., Zhang, M.J., Du, W.T., 2013b. The change of Ningchan River Glacier No.3 at Lenglongling, east Qilian Mountain, China. *Sciences in Cold and Arid Regions* 5 (6), 709.
- Liu, B.K., Du, Y.E., Hu, A.J., 2014. The dynamic of Qinghai lake area for remote sensing monitoring and the analysis of its driving force. *Advanced Materials Research* 971–973, 2162–2167.
- Ma, L.L., Tian, L.D., Pu, J.C., Wang, P.L., 2010. Recent area and ice volume change of Kangwure Glacier in the middle of Himalayas. *Chinese Science Bulletin* 55 (20), 2088–2096.
- Ma, N., Szilagyi, J., Niu, G.Y., Zhang, Y.S., Zhang, T., Wang, B.B., Wu, Y.H., 2016. Evaporation variability of Nam Co Lake in the Tibetan Plateau and its role in recent rapid lake expansion. *Journal of Hydrology* 537, 27–35.
- Molg, T., Grosshauser, M., Hemp, A., Hofer, M., Marzeion, B., 2012. Limited forcing of glacier loss through land-cover change on Kilimanjaro. *Nature Climate Change* 2 (4), 254–258.
- Molg, T., Maussion, F., Scherer, D., 2014. Mid-latitude westerlies as a driver of glacier variability in monsoonal High Asia. *Nature Climate Change* 4 (1), 68–73.
- Naidu, C.V., Durgalakshmi, K., Krishna, K.M., Rao, S.R., Satyanarayana, G.C., Lakshminarayana, P., Rao, L.M., 2009. Is summer monsoon rainfall decreasing over India in the global warming era? *Journal of Geophysical Research Atmospheres* 114 (D24), 144–153.
- Neckel, N., Kropacek, J., Bolch, T., Hochschild, V., 2014. Glacier mass changes on the Tibetan Plateau 2003–2009 derived from ICESat laser altimetry measurements. *Environmental Research Letters* 9 (1).
- Pan, B.T., Zhang, G.L., Wang, J., Cao, B., Geng, H.P., Wang, J., Zhang, C., Ji, Y.P., 2012. Glacier changes from 1966–2009 in the Gongga Mountains, on the south-eastern margin of the Qinghai–Tibetan Plateau and their climatic forcing. *Cryosphere* 6 (5), 1087–1101.
- Phan, V.H., Lindenbergh, R., Menenti, M., 2012. ICESat derived elevation changes of Tibetan lakes between 2003 and 2009. *International Journal of Applied Earth Observation and Geoinformation* 17, 12–22.
- Pu, J.C., Yao, T.D., Tian, L.D., 2006. Change of the Gurenhekou Glacier in Yangbajain Area, Nyainqntanglha Range. *Journal of Glaciology & Geocryology* 28 (6), 861–864.
- Pu, J.C., Yao, T.D., Yang, M.X., Tian, L.D., Wang, N.L., Ageta, Y., Fujita, K., 2008. Rapid decrease of mass balance observed in the Xiao (Lesser) Dongkemadi Glacier, in the central Tibetan Plateau. *Hydrological Processes* 22 (16), 2953–2958.
- Qin, J., Yang, K., Liang, S.L., Guo, X.F., 2009. The altitudinal dependence of recent rapid warming over the Tibetan Plateau. *Climatic Change* 97 (1), 321–327.
- Rangwala, I., Miller, J.R., Ming, X., 2009. Warming in the Tibetan Plateau: possible influences of the changes in surface water vapor. *Geophysical Research Letters* 36 (6), 295–311.
- Sakai, A., Fujita, K., Duan, K., Pu, J.C., Nakawo, M., Yao, T.D., 2005. Five decades of shrinkage of July 1st glacier, Qilian Shan, China. *Journal of Glaciology* 52 (176), 11–16.
- Scherer, D., Bookhagen, B., Strecker, M.R., 2011. Spatially variable response of Himalayan glaciers to climate change affected by debris cover. *Nature Geoscience* 4 (3), 156–159.
- Shahtahmassebi, A., Yu, Z.L., Wang, K., Xu, H.W., Deng, J.S., Li, J.D., Luo, R.S., Wu, J., Moore, N., 2012. Monitoring rapid urban expansion using a multi-temporal RGB-imperious surface model. *Journal of Zhejiang University-Science A* 13 (2), 146–158.
- Shang, Z.H., Gibb, M.J., Leiber, F., Ismail, M., Ding, L.M., Guo, X.S., Long, R.J., 2014. The sustainable development of grassland–livestock systems on the Tibetan plateau: problems, strategies and prospects. *Rangeland Journal* 36 (3), 267–296.
- Shangquan, D.H., Liu, S.Y., Ding, Y.J., Ding, L.F., Xiong, L.B., Cai, D.H., Li, G., Lu, A.X., Zhang, S.Q., Zhang, Y., 2006. Monitoring the glacier changes in the Muztag Ata and Konggur mountains, east Pamirs, based on Chinese Glacier Inventory and recent satellite imagery. *Annals of Glaciology* 43 (43), 79–85.
- Shao, Z., Meng, X., Zhu, D., Zheng, D., Qiao, Z., Yang, C., Han, J.E., Jia, Y., Meng, Q., Lü, R., 2008. Characteristics of the change of major lakes on the Qinghai–Tibet Plateau in the last 25 years. *Frontiers of Earth Science* 2 (3), 364–377.
- Shen, F., Kuang, D.B., 2003. Remote Sensing Investigation and Survey of Qinghai Lake in the Past 25 Years. *Journal of Lake Science* 15 (4), 289–296.
- Shen, G.Z., Liao, J.J., Guo, H.D., Li, Y.K., 2014. Study on the relationship between the lake area variations of Qinghai–Tibetan Plateau and the corresponding climate change in their basins. In: 35th International Symposium on Remote Sensing of Environment (Isrs35), pp. 17.
- Sheng, Z.P., Liu, Y.Q., Michelsen, A., Xu, D., 2005. Comparative study in water resources development of western regions in the US and China. *Transactions of the Asae* 48 (3), 1015–1024.
- Shi, P.H., Duan, K.Q., Liu, H.C., Yang, J.H., Zhang, X., Sun, J.Y., 2016. Response of Xiao Dongkemadi Glacier in the central Tibetan Plateau to the current climate change and future scenarios by 2050. *Journal of Mountain Science* 13 (1), 13–28.
- Song, C.Q., Huang, B., Ke, L.H., 2013. Modeling and analysis of lake water storage changes on the Tibetan Plateau using multi-mission satellite data. *Remote Sensing of Environment* 135 (4), 25–35.
- Song, C., Huang, B., Ke, L., 2014a. Modeling and analysis of lake water storage changes on the Tibetan Plateau using multi-mission satellite data. *Remote Sensing of Environment* 135 (4), 25–35.
- Song, C.Q., Huang, B., Ke, L.H., Richards, K.S., 2014b. Remote sensing of alpine lake water environment changes on the Tibetan Plateau and surroundings: A review. *Isprs Journal of Photogrammetry and Remote Sensing* 92, 26–37.
- Song, C.Q., Huang, B., Ke, L.H., Richards, K.S., 2014c. Seasonal and abrupt changes in the water level of closed lakes on the Tibetan Plateau and implications for climate impacts. *Journal of Hydrology* 514, 131–144.
- Song, C.Q., Huang, B., Richards, K., Ke, L.H., Phan, V.H., 2014d. Accelerated lake expansion on the Tibetan Plateau in the 2000s: Induced by glacial melting or other processes? *Water Resources Research* 50 (4), 3170–3186.
- Song, C.Q., Ke, L.H., Huang, B., Richards, K.S., 2015. Can mountain glacier melting explains the GRACE-observed mass loss in the southeast Tibetan Plateau: From a climate perspective? *Global and Planetary Change* 124, 1–9.
- Stofberg, S.F., Essink, G.H.P.O., Pauw, P.S., de Louw, P.G.B., Leijnse, A., van der Zee, S.E.A.T.M., 2017. Fresh water lens persistence and root zone salinization hazard under temperate climate. *Water Resources Management* 31 (2), 689–702.
- Sun, J., Cheng, G.W., Li, W.P., 2013a. Meta-analysis of relationships between environmental factors and aboveground biomass in the alpine grassland on the Tibetan Plateau. *Biogeosciences* 10 (3), 1707–1715.
- Sun, J., Cheng, G.W., Li, W.P., Sha, Y.K., Yang, Y.C., 2013b. On the variation of NDVI with the principal climatic elements in the Tibetan Plateau. *Remote Sensing* 5 (4), 1894–1911.
- Sun, J., Qin, X.J., Yang, J., 2016. The response of vegetation dynamics of the different alpine grassland types to temperature and precipitation on the Tibetan Plateau. *Environmental Monitoring and Assessment* 188 (1).
- Tang, X.L., Lv, X., He, Y., 2013. Features of climate change and their effects on glacier snow melting in Xinjiang, China. *Comptes Rendus Geoscience* 345 (2), 93–100.
- Thompson, L.G., Mountain, K., 2006. Abrupt tropical climate change: past and present. *Proceedings of the National Academy of Sciences* 103 (28), 10536–10543.
- Tian, L.D., Zong, J.B., Yao, T.D., Ma, L.L., Pu, J.C., Zhu, D.Y., 2014. Direct measurement of glacier thinning on the southern Tibetan Plateau (Gurenhekou, Kangwure and Naimona'nyi glaciers). *Journal of Glaciology* 60 (223), 879–888.
- Wagnon, P., Linda, A., Arnaud, Y., Kumar, R., Sharma, P., Vincent, C., Pottakkal, J.G., Berthier, E., Ramanathan, A., Hasnain, S.I., 2007. Four years of mass balance on Chhota Shigri Glacier, Himachal Pradesh, India, a new benchmark glacier in the western Himalaya. *Journal of Glaciology* 53 (183), 603–611.
- Wan, W., Xiao, P.F., Feng, X.Z., Li, H., Ma, R.H., Duan, H.T., Zhao, L.M., 2014. Monitoring lake changes of Qinghai–Tibetan Plateau over the past 30 years using satellite remote sensing data. *Chinese Science Bulletin* 59 (10), 1021–1035.
- Wang, B., Bao, Q., Hoskins, B., Wu, G.X., Liu, Y.M., 2008a. Tibetan Plateau warming and precipitation changes in East Asia. *Geophysical Research Letters* 35 (14).
- Wang, X., Xie, Z.C., Li, Q.Y., Wang, S.H., Cheng, L., 2008b. Sensitivity analysis of glacier systems to climate warming in China. *Journal of Geographical Sciences* 18 (2), 190–200.
- Wang, Y.T., Hou, S.G., Hong, S.M., Hur, S.D., Liu, Y.P., 2008c. Glacier extent and volume change (1966–2000) on the Su–Jo Mountain in northeastern Tibetan Plateau, China. *Journal of Mountain Science* 5 (4), 299–309.
- Wang, S.J., He, Y.Q., Song, X.D., 2010. Impacts of climate warming on alpine glacier tourism and adaptive measures: a case study of Baishui Glacier No. 1 in Yulong Snow

- Mountain, Southwestern China. *Journal of Earth Science* 21 (2), 166–178.
- Wang, L.P., Xie, Z.C., Liu, S.Y., Ding, L.F., Yin, J.Q., Li-Zong, W.U., 2011a. Glacierized area variation and its response to climate change in Qangtang Plateau during 1970–2000. *Journal of Glaciology & Geocryology* 33 (5), 979–990.
- Wang, X., Liu, S.Y., Mo, H.W., Yao, X.J., Jiang, Z.L., Guo, W.Q., 2011b. Expansion of glacial lakes and its implication for climate changes in the Chinese Himalaya. *Acta Geographica Sinica* 144 (2), 104–110.
- Wang, X.W., Gong, P., Zhao, Y.Y., Xu, Y., Cheng, X., Niu, Z.G., Luo, Z.C., Huang, H.B., Sun, F.D., Li, X.W., 2013a. Water-level changes in China's large lakes determined from ICESat/GLAS data. *Remote Sensing of Environment* 132, 131–144.
- Wang, X., Siegert, F., Zhou, A.G., Franke, J., 2013b. Glacier and glacial lake changes and their relationship in the context of climate change, Central Tibetan Plateau 1972–2010. *Global and Planetary Change* 111, 246–257.
- Wang, S.J., Du, J.K., He, Y.Q., 2014. Spatial-temporal characteristics of a temperate-glacier's active-layer temperature and its responses to climate change: a case study of Baishui Glacier No. 1, Southeastern Tibetan Plateau. *Journal of Earth Science* 25 (4), 727–734.
- Wang, J., Cui, Y.H., He, X.B., Zhang, J., Yan, S.J., 2015. Surface albedo variation and its influencing factors over dongkemadi glacier, Central Tibetan Plateau. *Advances in Meteorology* 2015 (1), 1–10.
- Wang, P.Y., Li, Z.Q., Yu, G.B., Li, H.L., Wang, W.B., Huai, B.J., Zhou, P., Jin, S., Wang, L., Zhang, H., 2016. Glacier shrinkage in the Daxue and Danghenan ranges of the western Qilian Mountains, China, from 1957 to 2010. *Environmental Earth Sciences* 75 (2).
- Wei, J.F., Liu, S.F., Guo, W.Q., Xu, J.L., Bao, W.J., Shangguan, D.H., 2015. Changes in Glacier Volume in the North Bank of the Bangong Co Basin from 1968 to 2007 Based on Historical Topographic Maps, SRTM, and ASTER Stereo Images. *Arctic, Antarctic, and Alpine Research* 47 (2), 301–311.
- Wu, Y.H., Zhu, L.P., 2008. The response of lake–glacier variations to climate change in Nam Co Catchment, central Tibetan Plateau, during 1970–2000. *Journal of Geographical Sciences* 18 (2), 177–189.
- Wu, Z.H., Jiang, Q.G., Cheng, B., 2007a. Study of dynamic changes of lakes in Qinghai–Tibet Plateau based on remote sensing and GIS. *Global Geology* 26 (01), 66–70.
- Wu, Y.H., Zhu, L.P., Ye, Q.H., Wang, L.B., 2007b. The Response of Lake–Glacier Area Change to Climate Variations in Namco Basin, Central Tibetan Plateau, during the Last Three Decades. *Acta Geographica Sinica* 62 (3), 301–311.
- Wu, Y.H., Zhang, H.X., Zhang, B., Chen, D.M., Lei, L.P., 2014. Long-Term Changes of Lake Level and Water Budget in the Nam Co Lake Basin, Central Tibetan Plateau. *Journal of Hydrometeorology* 15 (3), 1312–1322.
- Xie, Z.C., Wang, X., Feng, Q.H., Kang, E.S., Liu, C.H., Li, Q.Y., 2006. Modeling the response of glacier systems to climate warming in China. *Annals of Glaciology* 43 (1), 313–316.
- Xie, H., Ye, J.S., Liu, X.M., Chong, Y.E., 2010a. Warming and drying trends on the Tibetan Plateau (1971–2005). *Theoretical and Applied Climatology* 101 (3), 241–253.
- Xie, Z.C., Su, Z., Shen, Y.P., Feng, Q.H., 2010b. Mass balance and water exchange of Hailuoguo Glacier in Mount Gongga and their influence on Glacial Melt Runoff. *Journal of Glaciology & Geocryology* 23 (01), 7–15.
- Xing, Q., Li, Z., Zhou, J.M., 2010. Monitoring thickness change of the Dongkemadi Glacier on Qinghai–Tibetan Plateau Using Srtm Dem and map-based topographic data. *Ieee International Geoscience and Remote Sensing Symposium* 2010, 1769–1771.
- Xu, Z.X., Kong, T.L., Zhao, F.F., 2006. Analysis of climate change in Tibetan Plateau over the past 40 years. *Journal of Subtropical Resources & Environment* 1 (03), 24–32.
- Xu, Z.X., Gong, T.L., Li, J.Y., 2008. Decadal trend of climate in the Tibetan Plateau—regional temperature and precipitation. *Hydrological Processes* 22 (16), 3056–3065.
- Xu, B.Q., Cao, J.J., Hansen, J., Yao, T.D., Joswia, D.R., Wang, N.L., Wu, G.J., Mo, W., Zhao, H.B., Wei, Y., 2009. Black soot and the survival of Tibetan glaciers. *Proceedings of the National Academy of Sciences* 106 (52), 22114–22118.
- Yan, L.J., Zheng, M.P., 2015. The response of lake variations to climate change in the past forty years: A case study of the northeastern Tibetan Plateau and adjacent areas, China. *Quaternary International* 371, 31–48.
- Yang, W., Yao, T.D., Xu, B.Q., Wu, G.J., Ma, L.L., Xin, X.D., 2008a. Quick ice mass loss and abrupt retreat of the maritime glaciers in the Kangri Karpo Mountains, southeast Tibetan Plateau. *Chinese Science Bulletin* 53 (16), 2547–2551.
- Yang, Y.H., Fang, J.Y., Tang, Y.H., Ji, C.J., Zheng, C.Y., He, J.S., Zhu, B.A., 2008b. Storage, patterns and controls of soil organic carbon in the Tibetan grasslands. *Global Change Biology* 14 (7), 1592–1599.
- Yang, K., Ye, B.S., Zhou, D.G., Wu, B.Y., Foken, T., Qin, J., Zhou, Z.Y., 2011. Response of hydrological cycle to recent climate changes in the Tibetan Plateau. *Climatic Change* 109 (3), 517–534.
- Yang, J.P., Ding, Y.J., Liu, S.Y., Tan, C.P., 2015. Vulnerability of mountain glaciers in China to climate change. *Advances in Climate Change Research* 6 (3–4), 171–180.
- Yao, T.D., Pu, J.C., Lu, A.X., Wang, Y.Q., Yu, W.S., 2007. Recent Glacial Retreat and Its Impact on Hydrological Processes on the Tibetan Plateau, China, and Surrounding Regions. *Arctic Antarctic & Alpine Research* 39 (4), 642–650.
- Yao, T.D., Thompson, L., Yang, W., Yu, W.S., Gao, Y., Guo, X.J., Yang, X.X., Duan, K.Q., Zhao, H.B., Xu, B.Q., Pu, J.C., Lu, A.X., Xiang, Y., Kattel, D.B., Joswiak, D., 2012. Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings. *Nature Climate Change* 2 (9), 663–667.
- Yao, T.D., Masson-Delmotte, V., Gao, J., Yu, W.S., Yang, X.X., Risi, C., Sturm, C., Werner, M., Zhao, H.B., He, Y., Ren, W., Tian, L.D., Shi, C.M., Hou, S.G., 2013. A review of climatic controls on $\delta 18\text{O}$ in precipitation over the Tibetan Plateau: Observations and simulations. *Reviews of Geophysics* 51 (4), 525–548.
- Ye, Q.H., Kang, S.C., Chen, F., Wang, J.H., 2006. Monitoring glacier variations on Geladandong mountain, central Tibetan Plateau, from 1969 to 2002 using remote-sensing and GIS technologies. *Journal of Glaciology* 52 (179), 537–545.
- Ye, Q.H., Zhu, L.P., Zheng, H.X., Naruse, R.J., Zhang, X.Q., Kang, S.C., 2007. Glacier and lake variations in the Yamzhog Yumco basin, southern Tibetan Plateau, from 1980 to 2000 using remote-sensing and GIS technologies. *Journal of Glaciology* 53 (183), 673–676.
- Ye, Q.H., Yao, T.D., Zheng, H.X., Zhang, X.Q., 2008a. Glacier and lake co-variations and their responses to climate change in the Mapam Yumco Basin on Tibet. *Geographical Research* 27 (5), 1178–1190.
- Ye, Q.H., Yao, T.D., Naruse, R.J., 2008b. Glacier and lake variations in the Mapam Yumco basin, western Himalaya of the Tibetan Plateau, from 1974 to 2003 using remote-sensing and GIS technologies. *Journal of Glaciology* 54 (188), 933–935.
- Ye, Q.H., Zhong, Z.W., Kang, S.C., Stein, A., Wei, Q.F., Liu, J.S., 2009. Monitoring Glacier and Supra-glacier Lakes from Space in Mt. Qomolangma Region of the Himalayas on the Tibetan Plateau in China. *Journal of Mountain Science* 6 (3), 211–220.
- Yi, G.H., Zhang, T.B., 2015. Delayed response of lake area change to climate change in Siling Co Lake, Tibetan Plateau, from 2003 to 2013. *International Journal of Environmental Research and Public Health* 12 (11), 13886–13900.
- Yi, G.H., Deng, W., Li, A.N., Zhang, T.B., 2015. Response of lakes to climate change in Xainza basin Tibetan Plateau using multi-mission satellite data from 1976 to 2008. *Journal of Mountain Science* 12 (3), 604–613.
- You, Q., Kang, S., Flügel, W.A., Sanchez-Lorenzo, A., Yan, Y., Xu, Y., Huang, J., 2009. Does a weekend effect in diurnal temperature range exist in the eastern and central Tibetan Plateau? *Environmental Research Letters* 4 (4), 549–567.
- Yu, W.S., Yao, T.D., Kang, S.C., Pu, J.C., Yang, W., Gao, T.G., Zhao, H.B., Zhou, H., Li, S.H., Wang, W.C., Ma, L.L., 2013. Different region climate regimes and topography affect the changes in area and mass balance of glaciers on the north and south slopes of the same glacierized massif (the West Nyainqentanglha Range, Tibetan Plateau). *Journal of Hydrology* 495, 64–73.
- He, Y.Q., Yang, X.M., Jia, W.X., He, X.Z., Song, B., Zhang, N.N., Liu, Q., 2008. Changes of the Hailuoguo Glacier, Mt. Gongga, China, against the Background of Global Warming in the Last Several Decades. *Journal of China University of Geosciences* 19 (3), 271–281.
- Zhang, T., Ren, J.W., Kang, S.C., 2004. Lanong Glacier retreat in Nyainqentanglha Range of Tibetan Plateau during 1970–2003. *Journal of Glaciology & Geocryology* 26 (6), 736–739.
- Zhang, Y.Q., Liu, C.M., Tang, Y.H., Yang, Y.H., 2007. Trends in pan evaporation and reference and actual evapotranspiration across the Tibetan Plateau. *Journal of Geophysical Research* 112 (D12).
- Zhang, X., Ren, Y., Yin, Z.Y., Lin, Z., Zheng, D., 2009. Spatial and temporal variation patterns of reference evapotranspiration across the Qinghai–Tibetan Plateau during 1971–2004. *Journal of Geophysical Research* 114 (D15).
- Zhang, Y., Fujita, K.J., Liu, S.Y., Liu, Q., Wang, X., 2010. Multi-decadal ice-velocity and elevation changes of a monsoonal maritime glacier: Hailuoguo glacier, China. *Journal of Glaciology* 56 (195), 65–74.
- Zhang, G.Q., Xie, H.J., Kang, S.C., Yi, D.H., Ackley, S.F., 2011. Monitoring lake level changes on the Tibetan Plateau using ICESat altimetry data (2003–2009). *Remote Sensing of Environment* 115 (7), 1733–1742.
- Zhang, B., Wu, Y.H., Zhu, L.P., Wang, J.B., Li, J.S., Chen, D.M., 2011a. Estimation and trend detection of water storage at Nam Co Lake, central Tibetan Plateau. *Journal of Hydrology* 405 (1–2), 161–170.
- Zhang, G.Q., Xie, H.J., Duan, S.Q., Tian, M.Z., Yi, D.H., 2011b. Water level variation of Lake Qinghai from satellite and in situ measurements under climate change. *Journal of Applied Remote Sensing* 5 (1), 3532.
- Zhang, Y., Hirabayashi, Y., Liu, S.Y., 2012a. Catchment-scale reconstruction of glacier mass balance using observations and global climate data: Case study of the Hailuoguo catchment, south-eastern Tibetan Plateau. *Journal of Hydrology* 444, 146–160.
- Zhang, Y.S., Liu, S.Y., Shangguan, D.H., Li, J., Zhao, J.D., 2012b. Thinning and shrinkage of Laohuguo No. 12 glacier in the Western Qilian Mountains, China, from 1957 to 2007. *Journal of Mountain Science* 9 (3), 343–350.
- Zhang, M.J., Wang, S.J., Li, Z.Q., Wang, F.T., 2012c. Glacier area shrinkage in China and its climatic background during the past half century. *Journal of Geographical Sciences* 22 (1), 15–28.
- Zhang, G.Q., Yao, T.D., Xie, H.J., Kang, S.C., Lei, Y.B., 2013a. Increased mass over the Tibetan Plateau: From lakes or glaciers? *Geophysical Research Letters* 40 (10), 2125–2130.
- Zhang, J., Braaten, D., Li, X., She, J., Tao, F., 2013b. An inventory of glacier changes between 1973 and 2011 for the Geladandong Mountain area, China. *The Cryosphere Discussions* 7 (1), 507–531.
- Zhang, B., Wu, Y.H., Lei, L.P., Li, J.S., Liu, L.L., Chen, D.M., Wang, J.B., 2013c. Monitoring changes of snow cover, lake and vegetation phenology in Nam Co Lake Basin (Tibetan Plateau) using remote SENSING (2000–2009). *Journal of Great Lakes Research* 39 (2), 224–233.
- Zhang, G.Q., Xie, H.J., Yao, T.D., Kang, S.C., 2013d. Water balance estimates of ten greatest lakes in China using ICESat and Landsat data. *Chinese Science Bulletin* 58 (31), 3815–3829.
- Zhang, F., Shi, X., Zeng, C., 2015a. Study on Water Level Change of Yamzhog Yumco Lake in south Tibet. In: EGU General Assembly Conference.
- Zhang, G.Q., Yao, T.D., Xie, H.J., Wang, W.C., Yang, W., 2015b. An inventory of glacial lakes in the Third Pole region and their changes in response to global warming. *Global and Planetary Change* 131, 148–157.
- Zhang, Z., Xu, J.L., Liu, S.Y., Guo, W.Q., Wei, J.F., Feng, T., 2016. Glacier changes since the early 1960s, eastern Pamir, China. *Journal of Mountain Science* 13 (2), 276–291.
- Zhao, H.B., Xu, B.Q., Yao, T.D., Wu, G.J., Lin, S.B., Gao, J., Wang, M., 2012. Deuterium excess record in a southern Tibetan ice core and its potential climatic implications. *Climate Dynamics* 38 (9–10), 1791–1803.
- Zhisheng, A., Kutzbach, J.E., Prell, W.L., Porter, S.C., 2001. Evolution of Asian monsoons

- and phased uplift of the Himalaya–Tibetan plateau since Late Miocene times. *Nature* 411 (6833), 62–66.
- Zhou, S.Q., Kang, S.C., Chen, F., Joswiak, D.R., 2013. Water balance observations reveal significant subsurface water seepage from Lake Nam Co, south-central Tibetan Plateau. *Journal of Hydrology* 491, 89–99.
- Zhou, J., Wang, L., Zhang, Y.S., Guo, Y.H., Li, X.P., Liu, W.B., 2015. Exploring the water storage changes in the largest lake (Selin Co) over the Tibetan Plateau during 2003–2012 from a basin-wide hydrological modeling. *Water Resources Research* 51 (10), 8060–8086.
- Zhu, L.P., Xie, M.P., Wu, Y.H., 2010. Quantitative analysis of lake area variations and the influence factors from 1971 to 2004 in the Nam Co basin of the Tibetan Plateau. *Science Bulletin* 55 (13), 1294–1303.
- Li, Z.X., He, Y.Q., Pu, T., Jia, W.X., He, X.Z., Pang, H.X., Zhang, N.N., Liu, Q., Wang, S.J., Zhu, G.F., Wang, S.X., Chang, L., Du, J.K., Xin, H.J., 2010. Changes of climate, glaciers and runoff in China's monsoonal temperate glacier region during the last several decades. *Quaternary International* 218 (1–2), 13–28.